

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/55, C07D 243/14		A1	(11) International Publication Number: WO 96/00574
			(43) International Publication Date: 11 January 1996 (11.01.96)
(21) International Application Number: PCT/US95/08146		Schwenksville, PA 19473 (US). UZINSKAS, Irene, Nijole [US/US]; 154 Vassar Circle, Villanova, PA 19085 (US).	
(22) International Filing Date: 29 June 1995 (29.06.95)		(74) Agents: KINZIG, Charles, M. et al.; Smithkline Beecham Corporation, Corporate Intellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).	
(30) Priority Data: 08/267,695 29 June 1994 (29.06.94) US		(81) Designated States: JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(60) Parent Application or Grant (63) Related by Continuation US 08/267,695 (CIP) Filed on 29 June 1994 (29.06.94)		Published With international search report.	
(71) Applicant (for all designated States except US): SMITHKLINE BEECHAM CORPORATION [US/US]; Corporate Intellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): COUSINS, Russell, Donovan [US/US]; 2053 Kings Row, Oxford, PA 19363 (US). KEENAN, Richard, McCulloch [US/US]; 796 Bass Cove, Malvern, PA 19355 (US). KWON, Chet [US/US]; 649 South Henderson Road, King of Prussia, PA 19406 (US). MILLER, William, Henry [-/-]; 333 Fell Lane,			

(54) Title: VITRONECTIN RECEPTOR ANTAGONISTS

(57) Abstract


Compounds of formula (I) are disclosed which are inhibitors of the vitronectin receptor, wherein X-X' is NR¹-CH, NC(O)R³-CH, N=C, CR¹=C, CHR¹-CH, O-CH or S-CH; R¹ is H, C₁-alkyl or benzyl; R² is (CH₂)_nCO₂H; R³ is H, C₁-alkyl, Ar-C₀₋₆alkyl, Het-C₀₋₆alkyl, or C₃-cycloalkyl-C₀₋₆alkyl; R⁴ is W-U, Y-(CHR⁵)_m-U or Z-C(O); R⁵ and R⁶ are independently chosen from H, C₁-alkyl, Ar-C₀₋₆alkyl, Het-C₀₋₆alkyl and C₃-cycloalkyl-C₀₋₆alkyl; m is 1 or 2; n is 1 or 2; U is NR¹C(O), C(O)NR¹, CH=CH, C=C, CH₂-CH₂, O-CH₂, CH₂-O or CH₂OCONR¹; W is (a), (b) or (c); R^a is H, OH, NO₂, N(R⁶)₂, CON(R⁶)₂, CH₂N(R⁶)₂, or R⁶HN-C(=NH); Y is NH₂, NHR⁶, N(R⁶)₂, C(O)N(R⁶)₂, OH, -N-OR⁶, (d), (e) or (f); Z is (g) or (h); R^d is H, N(R¹)₂, C₁-alkyl, CON(R¹)₂, OH, OR¹, or Ar-C₀₋₆alkyl; R^e is H, C₁-alkyl, 2- or 3-pyridinyl, 1-, 2- or 3-piperidinyl, or 2- or 4-pyrimidinyl; and pharmaceutically acceptable salts thereof.




(a)




(b)



(c)




(d)




(e)



(f)



(g)



(h)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

TITLE

Vitronectin Receptor Antagonists

5

FIELD OF THE INVENTION

This invention relates to pharmaceutically active compounds which inhibit the vitronectin receptor and are useful for the treatment of osteoporosis.

BACKGROUND OF THE INVENTION

10

Mammalian bone is constantly undergoing bone remodeling, which is a dynamic process of bone resorption and bone formation. These processes are mediated by specialized cell types: bone formation is the result of the deposition of mineralized bone by osteoblast cells, and bone resorption is the result of the dissolution of bone matrix by osteoclast cells. Many bone diseases are brought about by an imbalance of bone formation relative to bone resorption. For instance, diseases such as osteoporosis are characterized by a net loss of bone matrix. Thus, agents which inhibit bone resorption are useful for the treatment of such diseases.

15

An activated osteoclast resorbs bone by attaching to the bone matrix, and secreting proteolytic enzymes, organic acids and protons into the sealed compartment formed between its cell membrane and the bone matrix. The acidic environment and proteolytic enzymes effect the dissolution of bone in the sealed compartment to create pits, or lacuna, in the bone surface, which are apparent when the osteoclast detaches from the bone.

20

Recent studies have indicated that the attachment of osteoclasts to the bone matrix is mediated through cell surface adhesion receptors called integrins. For instance, Davies, *et al.*, *J. Cell Biol.*, **1989**, 109, 1817, disclose that the osteoclast

25

- 2 -

functional antigen, which is implicated in the regulation of bone resorption, is biochemically related to the vitronectin receptor. The vitronectin receptor, or the $\alpha_v\beta_3$ integrin, is known to bind to bone matrix proteins, such as osteopontin, bone sialoprotein and thrombospondin, which contain the tri-peptide Arg-Gly-Asp (or RGD) motif. Thus, Horton, *et al.*, *Exp. Cell Res.* **1991**, *195*, 368, disclose that RGD-containing peptides and an anti-vitronectin receptor antibody (23C6) inhibit dentine resorption and cell spreading by osteoclasts. In addition, Sato, *et al.*, *J. Cell Biol.* **1990**, *111*, 1713 disclose that echistatin, a snake venom peptide which contains the RGD sequence, is a potent inhibitor of bone resorption in tissue culture, and inhibits attachment of osteoclasts to bone. Fisher, *et al.*, *Endocrinology* **1993**, *132*, 1411, has further shown that echistatin inhibits bone resorption *in vivo* in the rat. EP 528 587 and 528 586 report substituted phenyl derivatives which inhibit osteoclast mediated bone resorption.

Bondinell, *et al.*, in WO 93/00095 (PCT/US92/05463) and PCT/US93/12436 disclose that certain compounds which have a substituted 6-7 bicyclic ring system are useful for inhibiting the fibrinogen receptor, which is an integrin ($\alpha_{IIb}\beta_3$) protein found on platelets. Other 6-7 bicyclic ring systems which inhibit the fibrinogen receptor are disclosed by Blackburn *et al.* in WO 93/08174 (PCT/US92/08788). It has now been discovered that certain appropriately substituted 1,4-benzodiazepine, -benzothiazepine, -benzoxazepine and 2-benzazepine compounds are potent inhibitors of the vitronectin receptor. In particular, it has been discovered that certain such compounds are surprisingly more potent inhibitors of the vitronectin receptor than the fibrinogen receptor and consequently are more useful in the treatment of diseases such as osteoporosis, cancer, atherosclerosis and for inhibiting restenosis of an artery.

SUMMARY OF THE INVENTION

This invention comprises compounds of the formula (I) as described hereinafter, which have pharmacological activity for the inhibition of the vitronectin receptor and are useful in the treatment of osteoporosis.


This invention is also a pharmaceutical composition comprising a compound according to formula (I) and a pharmaceutically carrier.

This invention is also a method of treating diseases which are mediated by ligands which bind to the vitronectin receptor. In a particular aspect, the compounds of this invention are useful for treating osteoporosis, atherosclerosis, restenosis and cancer.

- 3 -

DETAILED DESCRIPTION

This invention comprises compounds of formula (I):



(I)





5

wherein

 $X-X'$ is NR^1-CH , $NC(O)R^3-CH$, $N=C$, $CR^1=C$, CHR^1-CH , $O-CH$ or $S-CH$; R^1 is H, C_{1-6} alkyl or benzyl; R^2 is $(CH_2)_nCO_2H$;10 R^3 is H, C_{1-6} alkyl, Ar- C_{0-6} alkyl, Het- C_{0-6} alkyl, or C_{3-6} cycloalkyl- C_{0-6} alkyl; R^4 is W-U, $Y-(CHR^5)_m-U$ or $Z-C(O)$; R^5 and R^6 are independently chosen from H, C_{1-6} alkyl, Ar- C_{0-6} alkyl, Het- C_{0-6} alkyl and C_{3-6} cycloalkyl- C_{0-6} alkyl;

m is 1 or 2;

15 n is 1 or 2;

U is $NR^1C(O)$, $C(O)NR^1$, $CH=CH$, $C\equiv C$, CH_2-CH_2 , $O-CH_2$, CH_2-O or CH_2OCONR^1 ; R^a is H, OH, NO_2 , $N(R^6)_2$, $CON(R^6)_2$, $CH_2N(R^6)_2$, or $R^6HN-C(=NH)$;20 Y is NH_2 , NHR^6 , $N(R^6)_2$, $C(O)N(R^6)_2$, OH, $=N-OR^6$, R^d is H, $N(R^1)_2$, C_{1-4} alkyl, $CON(R^1)_2$, OH, OR^1 , or Ar- C_{0-4} alkyl; R^e is H, C_{1-4} alkyl, Het- C_{0-4} alkyl or Ar- C_{0-4} alkyl;

25 and pharmaceutically acceptable salts thereof,

- 4 -



provided that R^3 is not phenylethyl when R^4 is (3-amidino)phenylaminocarbonyl and X-X' is NH-CH.


The compounds of formula (I) inhibit the binding of vitronectin and other RGD-containing peptides to the vitronectin ($\alpha_v\beta_3$) receptor. Inhibition of the vitronectin receptor on osteoclasts inhibits osteoclastic bone resorption and is useful in the treatment of diseases wherein bone resorption is associated with pathology, such as osteoporosis.

Suitably, X-X' is NH-CH or CH₂-CH.

Preferably, U is NR¹CO, CONR¹ or CH₂OCONR¹. More preferably U is NR¹CO.

Suitably R^a is hydroxy or amino. Preferably R^a is amino.

Suitably W is  .. Preferably, W is  or (3-amidino)phenyl.

Suitably Y is , OH or NHR⁶. Suitably R⁵ is H, phenyl or C₁₋₆alkyl. Suitably R⁶ is benzyl, 2-pyridinyl or H.

Suitably Z is 1-piperazinyl or 1-piperidinyl.

Suitably R⁴ is Y-(CH₂)_mNCH₃CO.

Suitably R^e is H or substituted or unsubstituted phenyl, benzyl, 2- or 3-pyridinyl, 2- or 4-pyrimidinyl or 1-, 2- or 3-piperidinyl. When Z is 1-piperazinyl, R^e is preferably 2- or 3-pyridinyl or 2- or 4-pyrimidinyl. When Z is piperidinyl, R^e is preferably 1-piperidinyl.

Preferably Y is NH₂ or pyridinyl.

Preferably n is 1.

Preferably R¹ is H, methyl or phenylethyl.

Representative of the novel compounds of this invention are the following:

(±)-7-[[[(6-Amino-2-pyridinyl)amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;

(±)-8-[[[(6-amino-2-pyridinyl)amino]carbonyl]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid;

(±)-7-[[[(6-amino-3-pyridinyl)amino]carbonyl]-3-oxo-4-(2-phenylethyl)-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;

(±)-1-Acetyl-7-[[[(6-amino-2-pyridinyl)amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;

- 5 -

- (±)-2-methyl-3-oxo-8-[[2-(pyridinyl)carbonyl]amino]-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid;
- (±)-8-[(benzyloxycarbonyl)amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid;
- 5 (±)-7-[[[3-(aminoiminomethyl)phenyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;
- (±)-7-[[[3-(aminocarbonyl)phenyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;
- (±)-4-methyl-3-oxo-7-[1-(4-phenylpiperazinyl)carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;
- 10 (±)-4-methyl-3-oxo-7-[(1-piperazinyl)carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;
- (±)-4-methyl-3-oxo-7-[1-[4-(1-piperidinyl)piperidinyl]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;
- (±)-4-methyl-3-oxo-7-[1-[4-(2-pyridinyl)piperazinyl]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;
- 15 (±)-4-methyl-3-oxo-7-[1-[4-(phenylmethyl)piperazinyl]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;
- (±)-4-methyl-3-oxo-7-[1-[4-(2-pyrimidinyl)piperazinyl]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;
- 20 8-[[2S-amino-3-phenylpropanoyl]amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4RS-acetic acid;
- (±)-8-[[[(2-hydroxy-2-phenyl)ethyl]methylamino]carbonyl]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid;
- 25 (±)-7-[[[N-(2-hydroxyethyl)-N-methyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;
- (±)-8-[[[(2-(2-pyridinylamino)acetyl)amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid;
- (±)-4-methyl-3-oxo-7-[[[(2-phenylaminoethyl)amino]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;
- 30 (±)-8-[(2-aminoacetyl)amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid;
- (+/-)-8-[(3-aminopropanoyl)amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid; and
- 35 (±)-4-methyl-3-oxo-7-[[[(3-pyridinyl)methyl]amino]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;

Other compounds useful in the method of inhibiting the vitronectin receptor are:

- 6 -

- (±)-7-[[[3-(Aminoiminomethyl)phenyl]amino]carbonyl]-3-oxo-4-(2-phenylethyl)-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid.
- (±)-7-[1-[4-(2-Methyl-4-pyridinyl)piperazinyl]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;
- 5 (±)-3-Oxo-4-(2-phenylethyl)-7-[1-[4-(4-pyridinyl)piperazinyl]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid; and
- (±)-4-Methyl-3-oxo-7-[1-[4-(4-pyridinyl)piperazinyl]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid.

Abbreviations and symbols commonly used in the peptide and chemical arts are used herein to describe the compounds of this invention. In general, the amino acid abbreviations follow the IUPAC-IUB Joint Commission on Biochemical Nomenclature as described in *Eur. J. Biochem.*, 158, 9 (1984).

C₁₋₄alkyl as applied herein means an optionally substituted alkyl group of 1 to 4 carbon atoms, and includes methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl and t-butyl. C₁₋₆alkyl additionally includes pentyl, n-pentyl, isopentyl, neopentyl and hexyl and the simple aliphatic isomers thereof. C₀₋₄alkyl and C₀₋₆alkyl additionally indicate that no alkyl group need be present (*e.g.*, that a covalent bond is present).

A substituent on a C₁₋₆ alkyl group, may be on any carbon atom which results in a stable structure, and is available by conventional synthetic techniques. Suitable substituents are C₁₋₄alkyl, OR¹, SR¹, C₁₋₄alkyl, C₁₋₄alkylsulfonyl, C₁₋₄alkylsulfoxyl, -CN, N(R¹)₂, CH₂N(R¹)₂, -NO₂, -CF₃, -CO₂R³, -CON(R¹)₂, -COR¹, -NR¹C(O)R¹, OH, F, Cl, Br, I, or CF₃S(O)_r, wherein r is 0 to 2.

Ar, or aryl, as applied herein, means phenyl or naphthyl, or phenyl or naphthyl substituted by one to three substituents, such as those defined above for alkyl, especially C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkthio, trifluoroalkyl, OH, F, Cl, Br, I or a 1,2-methylene dioxy group.

Het, or heterocycle, indicates an optionally substituted five or six membered monocyclic ring, or a nine or ten-membered bicyclic ring containing one to three heteroatoms chosen from the group of nitrogen, oxygen and sulfur, which are stable and available by conventional chemical synthesis. Illustrative heterocycles are benzofuryl, benzimidazole, benzopyran, benzothiophene, furan, imidazole, indoline, morpholine, piperidine, piperazine, pyrrole, pyrrolidine, tetrahydropyridine, pyridine, thiazole, thiophene, quinoline, isoquinoline, and tetra- and perhydro- quinoline and isoquinoline. Any accessible combination of up to three substituents on the Het ring, such as those defined above for alkyl that are

- 7 -

available by chemical synthesis and are stable are within the scope of this invention.

C₃₋₇cycloalkyl refers to an optionally substituted carbocyclic system of three to seven carbon atoms, which may contain up to two unsaturated carbon-carbon bonds. Typical of C₃₋₇cycloalkyl are cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl and cycloheptyl. Any combination of up to three substituents, such as those defined above for alkyl, on the cycloalkyl ring that is available by conventional chemical synthesis and is stable, is within the scope of this invention.


Certain radical groups are abbreviated herein. t-Bu refers to the tertiary butyl radical, Boc refers to the t-butyloxycarbonyl radical, Fmoc refers to the fluorenylmethoxycarbonyl radical, Ph refers to the phenyl radical, Cbz refers to the benzyloxycarbonyl radical, BrZ refers to the o-bromobenzyloxycarbonyl radical, ClZ refers to the o-chlorobenzyloxycarbonyl radical, Bn refers to the benzyl radical, 4-MBzl refers to the 4-methyl benzyl radical, Me refers to methyl, Et refers to ethyl, Ac refers to acetyl, Alk refers to C₁₋₄alkyl, Nph refers to 1- or 2-naphthyl and cHex refers to cyclohexyl. MeArg is N^α-methyl arginine. Tet refers to 5-tetrazolyl.

Certain reagents are abbreviated herein. DCC refers to dicyclohexylcarbodiimide, DMAP refers to dimethylaminopyridine, DIEA refers to diisopropylethylamine, EDC refers to N-ethyl-N'(dimethylaminopropyl)-carbodiimide. HOBt refers to 1-hydroxybenzotriazole, THF refers to tetrahydrofuran, DMF refers to dimethyl formamide, NBS refers to N-bromosuccinimide, Pd/C refers to a palladium on carbon catalyst, DPPA refers to diphenylphosphoryl azide, BOP refers to benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate, HF refers to hydrofluoric acid, TEA refers to triethylamine, TFA refers to trifluoroacetic acid, PCC refers to pyridinium chlorochromate.

Compounds of the formula (I) are prepared by the general methods described in Schemes I-III.

- 8 -

Scheme I




- 5 a) EDC, HOBT, (i-Pr)₂NEt, DMF, 2-aminopyridine; b) SOCl₂, reflux; c) 2-2-aminopyridine, pyridine, CH₂Cl₂; d) 1.0 N LiOH, aqueous THF; e) acidification.

Methyl (±)-7-carboxy-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (**I-1**), prepared as described by Bondinell, et al. (WO 93/00095), is converted to an activated form of the carboxylic acid using, for example, EDC and HOBT or SOCl₂, and the activated form is subsequently reacted with an appropriate amine to afford the corresponding amide **I-2**. Many additional methods for converting a carboxylic acid to an amide are known, and can be found in standard reference books, such as "Compendium of Organic Synthetic Methods", Vol. I - VI (published by Wiley-Interscience). The methyl ester of **I-2** is hydrolyzed using aqueous base, for example, aqueous LiOH in THF or aqueous NaOH in methanol, and the intermediate carboxylate salt is acidified with a suitable acid, for instance TFA or HCl, to afford the carboxylic acid **I-3**. Alternatively, the intermediate carboxylate salt can be isolated, if desired.

- 9 -

Scheme II




5 a) TFA, CH_2Cl_2 ; b) 1.0 N LiOH, aqueous THF.

If the amine partner contains a protecting group, the protecting group can be removed either prior or subsequently to the ester hydrolysis step, using methods suitable for selective deprotection of the specific protecting group employed. Such methods are described in Green, "Protective Groups in Organic Synthesis" (published by Wiley-Interscience). For example, if the amine partner contains a nitrogen group which is protected by a tert-butoxycarbonyl (BOC) group, such as in compound II-1 (prepared by the general methods described above), the BOC group is removed using acidic conditions, such as TFA in CH_2Cl_2 or HCl in dioxane, to afford the intermediate ammonium salt II-2. Subsequent ester hydrolysis followed by acidification is accomplished as described in Scheme I to afford II-3.

- 10 -


Scheme III



5 a) H₂, PtO₂, 1.0 N HCl, MeOH; b) 1.0 N LiOH, aqueous THF

Piperidine-containing compounds, such as **III-3**, can be prepared either from a suitably N-protected piperidine derivative, according to the methods described in Schemes I and II, or from a pyridine precursor, such as **III-1**. For example, the pyridine subunit of **III-1** can be reduced to the corresponding piperidine group by hydrogenation over a suitable catalyst, preferably PtO₂, in the presence of acid, such as HCl. The resulting piperidinium salt **III-2** is then converted to compound **III-3** by the methods described in Scheme I.

The core 6-7 fused ring system is prepared from compounds of the general formula (II):




- 11 -

wherein R^{10} is NHR^1 , CO_2H and synthetic equivalents thereof, X and X' are as defined for formula (I) and R^2 and R^3 are as defined in formula (I) with any reactive groups protected. Representative methods for preparing the substituted benzodiazepine nucleus are well known in the art, *e.g.*, Hynes, *et al.*, *J. Het. Chem.*, 1988, 25, 1173; Muller, *et al.*, *Helv. Chim. Acta.*, 1982, 65, 2118; Mori, *et al.*, *Heterocycles*, 1981, 16, 1491. Similarly, methods for preparing benzazepines, 1,4-benzothiazepines, 1,4-benzoxazepines and 1,4-benzodiazepines are known and are disclosed, for instance, in Bondinell, *et al.*, International Patent Application WO 93/00095.


Representative methods for preparing the benzodiazepine nucleus are given by Schemes IV - VI. A representative method for preparing a benzazepine nucleus is given by Scheme VII. A representative method for preparing a benzothiazepine is given by Scheme VIII. An benzoxazepine nucleus may be prepared in the same manner as Scheme VIII, except substituting a benzyl alcohol for a benzyl thiol.

Scheme IV




- 12 -

Scheme V



5


Scheme VI



10

- 13 -


Scheme VII



NBS = N-bromosuccinimide

- 14 -

Scheme VIII



5 The simple tri-substituted benzene starting materials are commercially available or prepared by routine methods well known in the art.

Coupling reagents as used herein denote reagents which may be used to form peptide bonds. Typical coupling methods employ carbodiimides, activated anhydrides and esters and acyl halides. Reagents such as EDC, DCC, DPPA, BOP reagent, HOBt, N-hydroxysuccinimide and oxalyl chloride are typical.

10 Coupling methods to form peptide bonds are generally well known to the art. The methods of peptide synthesis generally set forth by Bodansky *et al.*, THE PRACTICE OF PEPTIDE SYNTHESIS, Springer-Verlag, Berlin, 1984, Ali *et al.* in *J. Med. Chem.*, 29, 984 (1986) and *J. Med. Chem.*, 30, 2291 (1987) are generally illustrative of the technique and are incorporated herein by reference.

15 Typically, the amine or aniline is coupled via its free amino group to an appropriate carboxylic acid substrate using a suitable carbodiimide coupling agent, such as N,N'-dicyclohexyl carbodiimide (DCC), optionally in the presence of catalysts such as 1-hydroxybenzotriazole (HOBt) and dimethylamino pyridine (DMAP). Other methods, such as the formation of activated esters, anhydrides or

20

- 15 -

acid halides, of the free carboxyl of a suitably protected acid substrate, and subsequent reaction with the free amine of a suitably protected amine, optionally in the presence of a base, are also suitable. For example, a protected Boc-amino acid or Cbz-amidino benzoic acid is treated in an anhydrous solvent, such as methylene chloride or tetrahydrofuran(THF), in the presence of a base, such as N-methyl morpholine, DMAP or a trialkylamine, with isobutyl chloroformate to form the "activated anhydride", which is subsequently reacted with the free amine of a second protected amino acid or aniline.

Acid addition salts of the compounds are prepared in a standard manner in a suitable solvent from the parent compound and an excess of an acid, such as hydrochloric, hydrobromic, hydrofluoric, sulfuric, phosphoric, acetic, trifluoroacetic, maleic, succinic or methanesulfonic. Certain of the compounds form inner salts or zwitterions which may be acceptable. Cationic salts are prepared by treating the parent compound with an excess of an alkaline reagent, such as a hydroxide, carbonate or alkoxide, containing the appropriate cation; or with an appropriate organic amine. Cations such as Li^+ , Na^+ , K^+ , Ca^{++} , Mg^{++} and NH_4^+ are specific examples of cations present in pharmaceutically acceptable salts.

This invention also provides a pharmaceutical composition which comprises a compound according to formula (I) and a pharmaceutically acceptable carrier. Accordingly, the compounds of formula (I) may be used in the manufacture of a medicament. Pharmaceutical compositions of the compounds of formula (I) prepared as hereinbefore described may be formulated as solutions or lyophilized powders for parenteral administration. Powders may be reconstituted by addition of a suitable diluent or other pharmaceutically acceptable carrier prior to use. The liquid formulation may be a buffered, isotonic, aqueous solution. Examples of suitable diluents are normal isotonic saline solution, standard 5% dextrose in water or buffered sodium or ammonium acetate solution. Such formulation is especially suitable for parenteral administration, but may also be used for oral administration or contained in a metered dose inhaler or nebulizer for insufflation. It may be desirable to add excipients such as polyvinylpyrrolidone, gelatin, hydroxy cellulose, acacia, polyethylene glycol, mannitol, sodium chloride or sodium citrate.

Alternately, these compounds may be encapsulated, tableted or prepared in an emulsion or syrup for oral administration. Pharmaceutically acceptable solid or liquid carriers may be added to enhance or stabilize the composition, or to facilitate preparation of the composition. Solid carriers include starch, lactose, calcium sulfate dihydrate, terra alba, magnesium stearate or stearic acid, talc, pectin, acacia, agar or gelatin. Liquid carriers include syrup, peanut oil, olive oil, saline and

- 16 -

water. The carrier may also include a sustained release material such as glyceryl monostearate or glyceryl distearate, alone or with a wax. The amount of solid carrier varies but, preferably, will be between about 20 mg to about 1 g per dosage unit. The pharmaceutical preparations are made following the conventional
5 techniques of pharmacy involving milling, mixing, granulating, and compressing, when necessary, for tablet forms; or milling, mixing and filling for hard gelatin capsule forms. When a liquid carrier is used, the preparation will be in the form of a syrup, elixir, emulsion or an aqueous or non-aqueous suspension. Such a liquid formulation may be administered directly p.o. or filled into a soft gelatin capsule.

10 For rectal administration, the compounds of this invention may also be combined with excipients such as cocoa butter, glycerin, gelatin or polyethylene glycols and molded into a suppository.

The compounds described herein are antagonists of the vitronectin receptor, and are useful for treating diseases wherein the underlying pathology is attributable
15 to ligand or cell which interacts with the vitronectin receptor. For instance, these compounds are useful for the treatment of diseases wherein loss of the bone matrix creates pathology. Thus, the instant compounds are useful for the treatment of osteoporosis, hyperparathyroidism, Paget's disease, hypercalcemia of malignancy, osteolytic lesions produced by bone metastasis, bone loss due to immobilization or
20 sex hormone deficiency. The compounds of this invention are also believed to have utility as antitumor, antiinflammatory, anti-angiogenic and anti-metastatic agents, and be useful in the treatment of cancer, atherosclerosis and restenosis.

The peptide is administered either orally or parenterally to the patient, in a manner such that the concentration of drug is sufficient to inhibit bone resorption,
25 or other such indication. The pharmaceutical composition containing the peptide is administered at an oral dose of between about 0.1 to about 50 mg/kg in a manner consistent with the condition of the patient. Preferably the oral dose would be about 0.5 to about 20 mg/kg. For acute therapy, parenteral administration is preferred. An intravenous infusion of the peptide in 5% dextrose in water or
30 normal saline, or a similar formulation with suitable excipients, is most effective, although an intramuscular bolus injection is also useful. Typically, the parenteral dose will be about 0.01 to about 100 mg/kg; preferably between 0.1 and 20 mg/kg. The compounds are administered one to four times daily at a level to achieve a total daily dose of about 0.4 to about 400 mg/kg/day. The precise level and method by
35 which the compounds are administered is readily determined by one routinely skilled in the art by comparing the blood level of the agent to the concentration required to have a therapeutic effect.

- 17 -

The compounds may be tested in one of several biological assays to determine the concentration of compound which is required to have a given pharmacological effect.

5 INHIBITION OF VITRONECTIN BINDING

Solid-Phase [³H]-SK&F-107260 Binding to $\alpha_v\beta_3$: Human placenta or human platelet $\alpha_v\beta_3$ (0.1-0.3 mg/mL) in buffer T (containing 2 mM CaCl₂ and 1% octylglucoside) was diluted with buffer T containing 1 mM CaCl₂, 1 mM MnCl₂, 1 mM MgCl₂ (buffer A) and 0.05% NaN₃, and then immediately added to 96-well ELISA plates (Corning, New York, NY) at 0.1 mL per well. 0.1 - 0.2 μ g of $\alpha_v\beta_3$ was added per well. The plates were incubated overnight at 4°C. At the time of the experiment, the wells were washed once with buffer A and were incubated with 0.1 mL of 3.5% bovine serum albumin in the same buffer for 1 hr at room temperature. Following incubation the wells were aspirated completely and washed twice with 0.2 mL buffer A.

Compounds were dissolved in 100% DMSO to give a 2 mM stock solution, which was diluted with binding buffer (15 mM Tris-HCl (pH 7.4), 100 mM NaCl, 1 mM CaCl₂, 1 mM MnCl₂, 1 mM MgCl₂) to a final compound concentration of 100 μ M. This solution is then diluted to the required final compound concentration. Various concentrations of unlabeled antagonists (0.001 - 100 μ M) were added to the wells in triplicates, followed by the addition of 5.0 nM of [³H]-SK&F-107260 (65 - 86 Ci/mmol).

The plates were incubated for 1 hr at room temperature. Following incubation the wells were aspirated completely and washed once with 0.2 mL of ice cold buffer A in a well-to-well fashion. The receptors were solubilized with 0.1 mL of 1% SDS and the bound [³H]-SK&F-107260 was determined by liquid scintillation counting with the addition of 3 mL Ready Safe in a Beckman LS Liquid Scintillation Counter, with 40% efficiency. Nonspecific binding of [³H]-SK&F-107260 was determined in the presence of 2 μ M SK&F-107260 and was consistently less than 1% of total radioligand input. The IC₅₀ (concentration of the antagonist to inhibit 50% binding of [³H]-SK&F-107260) was determined by a nonlinear, least squares curve-fitting routine, which was modified from the LUNDON-2 program. The K_i (dissociation constant of the antagonist) was calculated according to the equation: $K_i = IC_{50}/(1 + L/K_d)$, where L and K_d were the concentration and the dissociation constant of [³H]-SK&F-107260, respectively.

- 18 -

Compounds of the present invention inhibit vitronectin binding to SK&F 107260 in the concentration range of 0.1 to 25 micromolar. Preferred compounds inhibit vitronectin binding at a concentration of less than 1 micromolar.

Compounds of this invention are also tested for *in vitro* and *in vivo* bone resorption in assays standard in the art for evaluating inhibition of bone formation, such as the pit formation assay disclosed in EP 528 587, which may also be performed using human osteoclasts in place of rat osteoclasts, and the ovariectomized rat model, described by Wronski *et al.*, *Cells and Materials* 1991, Sup. 1, 69-74.

10

PARATHYROIDECTOMIZED RAT MODEL

Each experimental group consists of 5-6 male Sprague-Dawley rats. The rats are parathyroidectomized (by the vendor, Taconic Farms) 7 days prior to use. Twenty four hours prior to use, circulating ionized calcium levels are measured in whole blood immediately after it has been withdrawn by tail venipuncture into heparinized tubes. Rats are included if ionized Ca level (measured with a Ciba-Corning model 634 calcium pH analyzer) is ≤ 1.2 mM/L. The rats are then put on a diet of calcium-free chow and deionized water. At the start of the experiment the rats weigh approximately 100g. Baseline Ca levels are measured and the rats are administered control vehicle (saline) or compound (dissolved in saline) as a single intravenous (tail vein) bolus injection followed immediately by a single subcutaneous injection of either human parathyroid hormone 1-34 peptide (hPTH1-34, dose 0.2mg/kg in saline/0.1% bovine serum albumen, Bachem, Ca) or the PTH vehicle. The calcemic response to PTH (and any effect of compound on this response) is measured 2h after compound/PTH administration.

25

RAT ULNA DRIFT MODEL

Each experimental group consists of 8-10 male Sprague-Dawley or Wistar rats of approximately 30-40g body weight at the start of the experiment. The agent being tested is administered by an appropriate route as single or multiple daily doses for a period of seven days. Prior to administration of the first dose, the rats are given a single dose of a fluorescent marker (tetracycline 25mg/kg, or calcein 10mg/kg) that labels the position of bone forming surfaces at that point in time. After dosing of compound has been completed, the rats are killed and both forelimbs are removed at the elbow, the foot is removed at the ankle and the skin removed. The sample is frozen and mounted vertically on a microtome chuck. Cross sections of the midshaft region of the ulna are cut in the cryostat. The rate of bone resorption is measured morphometrically in the medial-dorsal portion of the cortical bone. The measurement is done as follows: the amount of bone

35

- 19 -

- resorbed at the periosteal surface is equal to the distance by which the periosteal surface has advanced towards the fluorescent label which had been incorporated at the endosteal bone formation surface on day zero; this distance is calculated by subtracting the width of bone between the label and the periosteal surface on day 7 from the width on day zero;
- 5 the resorption rate in microns per day is calculated by dividing the result by 7.

HUMAN OSTEOCLAST RESORPTION ASSAY ("PIT ASSAY")

- Aliquots of osteoclastoma-derived cell suspensions are removed from liquid nitrogen storage, warmed rapidly at 37°C and washed x1 in RPMI-1640 medium
- 10 by centrifugation (1000rpm, 5 mins at 4°C).
- Aspirate the medium and replace it with murine anti-HLA-DR antibody, diluted 1:3 in RPMI-1640 medium. Incubate for 30 mins on ice and mix the cell suspension frequently.
- The cells are washed x2 with cold RPMI-1640 by centrifugation (1000rpm, 5
- 15 mins at 4°C) and the cells are transferred to a sterile 15 ml centrifuge tube. The number of mononuclear cells are enumerated in an improved Neubauer counting chamber.
- Sufficient magnetic beads (5 / mononuclear cell), coated with goat anti-mouse IgG, are removed from their stock bottle and placed into 5 ml of fresh medium
- 20 (this washes away the toxic azide preservative). The medium is removed by immobilizing the beads on a magnet and is replaced with fresh medium.
- The beads are mixed with the cells and the suspension is incubated for 30 mins on ice. The suspension is mixed frequently.
- The bead-coated cells are immobilized on a magnet and the remaining cells
- 25 (osteoclast-rich fraction) are decanted into a sterile 50 ml centrifuge tube.
- Fresh medium is added to the bead-coated cells to dislodge any trapped osteoclasts. This wash process is repeated x10. The bead-coated cells are discarded.
- The osteoclasts are enumerated in a counting chamber, using a large-bore disposable plastic pasteur to charge the chamber with the sample.
- 30 • The cells are pelleted by centrifugation and the density of osteoclasts adjusted to 1.5×10^4 /ml in EMEM medium, supplemented with 10% fetal calf serum and 1.7g/litre of sodium bicarbonate.
- 3ml aliquots of the cell suspension (per treatment) are decanted into 15ml centrifuge tubes. The cells are pelleted by centrifugation.
- 35 • To each tube 3ml of the appropriate treatment are added (diluted to 50 uM in the EMEM medium). Also included are appropriate vehicle controls, a positive control

- 20 -

(87MEM1 diluted to 100 ug/ml) and an isotype control (IgG2a diluted to 100 ug/ml). Incubate at 37°C for 30 mins.

- 0.5ml aliquots of the cells are seeded onto sterile dentine slices in a 48-well plate and incubated at 37°C for 2 hours. Each treatment is screened in quadruplicate.
 - 5 • The slices are washed in six changes of warm PBS (10 ml / well in a 6-well plate) and then placed into fresh treatment or control. Incubate at 37°C for 48 hours.
- tartrate resistant acid phosphatase (trap) procedure** (selective stain for cells of the osteoclast lineage).
- The slices are washed in phosphate buffered saline and fixed in 2%
 - 10 gluteraldehyde (in 0.2M sodium cacodylate) for 5 mins.
 - They are washed in water and incubated in TRAP buffer for 5 mins at 37°C.
 - Following a wash in cold water they are incubated in cold acetate buffer / fast red garnet for 5 mins at 4°C.
 - Excess buffer is aspirated, and the slices are air dried following a wash in water.
 - 15 • The TRAP positive osteoclasts are enumerated by bright-field microscopy and are then removed from the surface of the dentine by sonication.
 - Pit volumes are determined using the Nikon/Lasertec ILM21W confocal microscope.

20 INHIBITION OF RGD-MEDIATED GPIIB-IIIa BINDING

Purification of GPIIb-IIIa

- Ten units of outdated, washed human platelets (obtained from Red Cross) were lyzed by gentle stirring in 3% octylglucoside, 20 mM Tris-HCl, pH 7.4, 140 mM NaCl, 2 mM CaCl₂ at 4°C for 2 h. The lysate was centrifuged at 100,000g for
- 25 1 h. The supernatant obtained was applied to a 5 mL lentil lectin sepharose 4B column (E.Y. Labs) preequilibrated with 20 mM Tris-HCl, pH 7.4, 100 mM NaCl, 2 mM CaCl₂, 1% octylglucoside (buffer A). After 2 h incubation, the column was washed with 50 mL cold buffer A. The lectin-retained GPIIb-IIIa was eluted with buffer A containing 10% dextrose. All procedures were performed at 4°C. The
- 30 GPIIb-IIIa obtained was >95% pure as shown by SDS polyacrylamide gel electrophoresis.

Incorporation of GPIIb-IIIa in Liposomes.

- A mixture of phosphatidylserine (70%) and phosphatidylcholine (30%)
- 35 (Avanti Polar Lipids) were dried to the walls of a glass tube under a stream of nitrogen. Purified GPIIb-IIIa was diluted to a final concentration of 0.5 mg/mL and mixed with the phospholipids in a protein:phospholipid ratio of 1:3 (w:w). The

- 21 -

- mixture was resuspended and sonicated in a bath sonicator for 5 min. The mixture was then dialyzed overnight using 12,000-14,000 molecular weight cutoff dialysis tubing against a 1000-fold excess of 50 mM Tris-HCl, pH 7.4, 100 mM NaCl, 2 mM CaCl₂ (with 2 changes). The GPIIb-IIIa-containing liposomes were centrifuged at 12,000g for 15 min and resuspended in the dialysis buffer at a final protein concentration of approximately 1 mg/mL. The liposomes were stored at -70C until needed.


Competitive Binding to GPIIb-IIIa

- The binding to the fibrinogen receptor (GPIIb-IIIa) was assayed by an indirect competitive binding method using [³H]-SK&F-107260 as an RGD-type ligand. The binding assay was performed in a 96-well filtration plate assembly (Millipore Corporation, Bedford, MA) using 0.22 μ m hydrophilic durapore membranes. The wells were precoated with 0.2 mL of 10 μ g/mL polylysine (Sigma Chemical Co., St. Louis, MO.) at room temperature for 1 h to block nonspecific binding. Various concentrations of unlabeled benzadiazapines were added to the wells in quadruplicate. [³H]-SK&F-107260 was applied to each well at a final concentration of 4.5 nM, followed by the addition of 1 μ g of the purified platelet GPIIb-IIIa-containing liposomes. The mixtures were incubated for 1 h at room temperature. The GPIIb-IIIa-bound [³H]-SK&F-107260 was separated from the unbound by filtration using a Millipore filtration manifold, followed by washing with ice-cold buffer (2 times, each 0.2 mL). Bound radioactivity remaining on the filters was counted in 1.5 mL Ready Solve (Beckman Instruments, Fullerton, CA) in a Beckman Liquid Scintillation Counter (Model LS6800), with 40% efficiency. Nonspecific binding was determined in the presence of 2 μ M unlabeled SK&F-107260 and was consistently less than 0.14% of the total radioactivity added to the samples. All data points are the mean of quadruplicate determinations.

- Competition binding data were analyzed by a nonlinear least-squares curve fitting procedure. This method provides the IC₅₀ of the antagonists (concentration of the antagonist which inhibits specific binding of [³H]-SK&F-107260 by 50% at equilibrium). The IC₅₀ is related to the equilibrium dissociation constant (K_i) of the antagonist based on the Cheng and Prusoff equation: $K_i = IC_{50}/(1+L/K_d)$, where L is the concentration of [³H]-SK&F-107260 used in the competitive binding assay (4.5 nM), and K_d is the dissociation constant of [³H]-SK&F-107260 which is 4.5 nM as determined by Scatchard analysis.

Preferred compounds of this invention have an affinity for the vitronectin receptor relative to the fibrinogen receptor of greater than 3:1. More preferred

compounds have a ratio of activity of greater than 10:1. The comparative results of the enhanced binding of the compounds of this invention to the vitronectin receptor relative to the fibrinogen receptor are given in Table 1 below:












5

Ex	R ⁴ = W-U		W	$\alpha_V\beta_3$ (μ M)	$\alpha_{IIb}\beta_3$ (μ M)	ratio*
	X ¹	U				
1	NH	NHCO		0.15	160	1067
2	CH ₂	NHCO		0.65	>50	77
3	NH	NHCO		9.18	110	12
4	NAc	NHCO		15.1	>100	>7
5	CH ₂	CONH		15.3	>50	>3
6	CH ₂	CH ₂ OCONH		7.50	17.5	2.3
7	NH	NHCO		1.50	5.05	3.4
8	NH	NHCO		12.3	>50	>4

* enhanced binding to vitronectin receptor vs fibrinogen receptor = $\alpha_{IIb}\beta_3/\alpha_V\beta_3$

- 23 -

R ⁴ =Z-CO						
Ex	X ¹		Z	$\alpha\gamma\beta_3$ (μM)	$\alpha\pi\beta_3$ (μM)	ratio*
9	NH	CO		9.23	>100	>11
10	NH	CO		1.10	2.13	>1.9
11	NH	CO		1.45	20.1	14
12	NH	CO		1.25	40.5	32
13	NH	CO		9.5	15.4	1.62
14	NH	CO		6.2	>50	>8

R ⁴ = Y-(CHR) _m U						
Ex	X ¹	-(CHR) _m -U	Y	$\frac{\alpha\gamma\beta_3}{(\mu M)}$	$\frac{\alpha\pi\beta_3}{(\mu M)}$	ratio*
15	CH ₂	CHPhCONH	NH ₂	3.90	37.5	9.6
16	CH ₂	CHPhCONMe	OH	9.50	28.0	2.9
17	NH	NMeCO		15.2	>50	>3
18	CH ₂	CH ₂ CONH		1.3	30.0	23
19	NH	CH ₂ CH ₂ NHCO	NHPh	22.0	>50	>2.3
20	CH ₂	CH ₂ CONH	NH ₂	1.55	45.0	29
21	CH ₂	CH ₂ CH ₂ CONH	NH ₂	0.85	52.0	61
22	NH	CH ₂ NHCO		9.02	27.0	3

Vascular smooth muscle cell migration assay

The compounds of the instant invention were tested for their ability to
 5 inhibit the migration and proliferation of smooth muscle tissue in an artery or vein

- 24 -

in order to assess their ability to prevent restenosis of an artery, such as that which typically occurs following angioplasty.

Rat or human aortic smooth muscle cells were used. The cell migration was monitored in a Transwell cell culture chamber by using a polycarbonate membrane with pores of 8 μm (Costar). The lower surface of the filter was coated with vitronectin. Cells were suspended in DMEM supplemented with 0.2% bovine serum albumin at a concentration of $2.5 - 5.0 \times 10^6$ cells/mL, and were pretreated with test compound at various concentrations for 20 min at 20°C. The solvent alone was used as control. 0.2 mL of the cell suspension was placed in the upper compartment of the chamber. The lower compartment contained 0.6 mL of DMEM supplemented with 0.2% bovine serum albumin. Incubation was carried out at 37°C in an atmosphere of 95% air/5% CO₂ for 24 hr. After incubation, the non-migrated cells on the upper surface of the filter were removed by gentle scraping. The filter was then fixed in methanol and stained with 10% Giemsa stain. Migration was measured either by a) counting the number of cells that had migrated to the lower surface of the filter or by b) extracting the stained cells with 10% acetic acid followed by determining the absorbance at 600 nm.

20

Examples

Nuclear magnetic resonance spectra were recorded at either 250 or 400 MHz using, respectively, a Bruker AM 250 or Bruker AC 400 spectrometer. CDCl₃ is deuteriochloroform, DMSO-d₆ is hexadeuteriodimethylsulfoxide, and CD₃OD is tetradeuteriomethanol. Chemical shifts are reported in parts per million (δ) downfield from the internal standard tetramethylsilane. Abbreviations for NMR data are as follows: s=singlet, d=doublet, t=triplet, q=quartet, m=multiplet, dd=doublet of doublets, dt=doublet of triplets, app=apparent, br=broad. J indicates the NMR coupling constant measured in Hertz. Infrared (IR) spectra were recorded on a Perkin-Elmer 683 infrared spectrometer in transmission mode. IR band positions are reported in inverse wavenumbers (cm⁻¹). Mass spectra were taken on either VG 70 FE, PE Syx API III, or VG ZAB HF instruments, using fast atom bombardment (FAB) or electrospray (ES) ionization techniques. Elemental analyses were obtained using a Perkin-Elmer 240C elemental analyzer. Melting points were taken on a Thomas-Hoover melting point apparatus and are uncorrected. All temperatures are reported in degrees Celsius.

35

Analtech Silica Gel GF and E. Merck Silica Gel 60 F-254 thin layer plates were used for thin layer chromatography. Both flash and gravity chromatography

- 25 -

were carried out on E. Merck Kieselgel 60 (230-400 mesh) silica gel. Analytical and preparative HPLC were carried out on Rainin or Beckman chromatographs. ODS refers to an octadecylsilyl derivatized silica gel chromatographic support. 5 μ Apex-ODS indicates an octadecylsilyl derivatized silica gel chromatographic support having a nominal particle size of 5 μ , made by Jones Chromatography, Littleton, Colorado. YMC ODS-AQ® is an ODS chromatographic support and is a registered trademark of YMC Co. Ltd., Kyoto, Japan. PRP-1® is a polymeric (styrene-divinylbenzene) chromatographic support, and is a registered trademark of Hamilton Co., Reno, Nevada. Celite® is a filter aid composed of acid-washed diatomaceous silica, and is a registered trademark of Manville Corp., Denver, Colorado.

Methyl (\pm)-7-carboxy-3-oxo-2-(2-phenylethyl)-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate, methyl (\pm)-7-carboxy-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate, and 3-amino-(N-benzyloxycarbonyl)benzamidine were prepared by the method of Bondinell, et al., WO 93/00095. Methyl (\pm)-8-carboxy-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetate was prepared by the method disclosed in Example 1 of WO 94/14776, except substituting 4-bromo-3-methylbenzoic acid for 3-bromo-4-methylbenzoic acid and substituting methylamine for phenethylamine therein. 4-(2-Methylpyridyl) piperazine was prepared by the procedure of Cross and Dickinson, US 4,806,536 (Feb. 21, 1989).

Example 1

25 Preparation of (\pm)-7-[[6-Amino-2-pyridinyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

a) Methyl (\pm)-7-[[6-amino-2-pyridinyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate

30 Diisopropylethylamine (0.29 g, 2.25 mmol) was added in one portion to a stirred mixture of 2,6-diaminopyridine (0.2 g, 1.8 mmol), methyl (\pm)-7-carboxy-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (0.44 g, 1.50 mmol), EDC (0.34 g, 1.8 mmol) and HOBt-H₂O (0.24 g, 1.8 mmol) in DMF (8 mL) at RT under argon. After 24 h, the brownish solution was poured into a mixture of ice-water (90 g) and 5% NaHCO₃ (10 mL). The resulting precipitate

35 was filtered and air-dried to give an off-white solid (0.53 g). Flash chromatography (silica gel, 7% MeOH/CH₂Cl₂) yielded the title compound (0.04 g, 7%): ¹H NMR

- 26 -

(250 MHz, CDCl₃) δ 6.26-8.20 (m, 6H), 5.47 (d, J=16 Hz, 1H), 5.12 (dd, J=11, 6 Hz, 1H), 4.59 (d, J=4 Hz, 2H), 3.80 (d, J=16 Hz, 1H), 3.76 (s, 3H), 3.09 (s, 3H), 3.00 (dd, J=16, 6 Hz, 1H), 2.68 (dd, J=16, 6 Hz, 1H); MS (ES) m/e 384.2 (M+H)⁺.

- 5 b) (\pm)-7-[[[(6-Amino-2-pyridinyl)amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

1.0 N LiOH (0.26 mL, 0.26 mmol) was added dropwise in RT to a mixture of methyl (\pm)-7-[[[(6-amino-2-pyridinyl)amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (0.044 g, 0.12 mmol) in THF (4 mL) and H₂O (5 mL). The resulting light yellowish-brown solution was stirred for 23 h, then was concentrated on the rotavap. The resulting residue was lyophilized to give the crude product as a yellowish powder (0.051 g). Preparative HPLC (PRP-1® column, step gradient, 10-15% CH₃CN/H₂O-0.1% TFA) afforded the title compound: ¹H NMR (250 MHz, DMSO-d₆) δ 10.75 (br, 1H), 6.46-8.29 (m, 6H),
15 5.73 (s, 1H), 5.53 (d, J=17 Hz, 1H), 5.17 (br, 1H), 3.88 (d, J=17 Hz, 1H), 2.96 (s, 3H), 2.83 (dd, J=17, 9 Hz, 1H), 2.58 (dd, J=17, 5 Hz, 1H); MS (ES) m/e 370.2 (M+H)⁺. Anal. Calcd for C₁₈H₁₉N₅O₄ · 9/4 CF₃CO₂H · 1/2 H₂O: C, 42.56; H, 3.53; N, 11.03. Found: C, 42.20; H, 3.02; N, 11.36.

20

Example 2

Preparation of (\pm)-8-[[[(6-amino-2-pyridinyl)amino]carbonyl]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid

- 25 a) Methyl (\pm)-8-[[[(6-amino-2-pyridinyl)amino]carbonyl]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetate

A mixture of methyl (\pm)-8-carboxy-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetate (300 mg, 1.03 mmol) and thionyl chloride (10 mL) was refluxed for 1 h. The resulting solution was concentrated to dryness to yield a
30 foam. This was dissolved in CH₂Cl₂ (10 mL) and added dropwise to a solution of 2,6-diaminopyridine (0.34 g, 3.09 mmol) in CH₂Cl₂ (10 mL) and DMF (3 mL) at RT under argon. After 110 min, the reaction mixture was partitioned between CH₂Cl₂ (150 mL) and 5% NaHCO₃ (70 mL). The organic layer was separated, washed sequentially with 5% NaHCO₃ (50 mL) and H₂O (50 mL), dried
35 (Na₂SO₄), and concentrated to leave a yellowish solid (0.69 g). Recrystallization from MeOH/CH₂Cl₂/EtOAc afforded the title compound (0.16 g, 41%): ¹H NMR (250 MHz, CDCl₃/DMSO-d₆) δ 6.63-10.00 (m, 6H), 5.71 (s, 2H), 5.30 (d, J=16

- 27 -

Hz, 1H), 4.07 (d, J=17 Hz, 1H), 3.83 (m, 1H), 3.61 (s, 3H), 2.93 (s, 3H), 2.71-3.25 (m, 3H), 2.45 (dd, J=17, 5 Hz, 1H); MS (ES) m/e 383.2 (M+H)⁺.

- 5 b) (±)-8-[[[(6-Amino-2-pyridinyl)amino]carbonyl]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid

1.0 N LiOH (0.92 mL, 0.92 mmol) was added dropwise at RT to a mixture of the compound of Example 2(a) (160 mg, 0.42 mmol) in THF (6.5 mL) and H₂O (8 mL). The resulting solution was stirred for 21.5 h, then concentrated on the rotavap. The resulting residue was lyophilized to give the crude product, and
10 purified by preparative HPLC (PRP-1® column, step gradient, 10-20% CH₃CN/H₂O-0.1% TFA) to yield the title compound: ¹H NMR (250 MHz, DMSO-d₆) δ 11.13 (br, 1H), 6.52-7.88 (m, 6H), 5.35 (d, J=16 Hz, 1H), 4.10 (d, J=17 Hz, 1H), 3.82 (m, 1H), 2.94 (s, 3H), 2.53-3.22 (m, 3H), 2.37 (dd, J=17, 5 Hz, 1H); MS (ES) m/e 369.2 (M+H)⁺. Anal. Calcd for C₁₉H₂₀N₄O₄ · 3/2 CF₃CO₂H:
15 C, 48.99; H, 4.02; N, 10.39. Found: C, 48.75; H, 4.23; N, 10.35.

Example 3

- 20 Preparation of (±)-7-[[[(6-amino-3-pyridinyl)amino]carbonyl]-3-oxo-4-(2-phenylethyl)-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

- a) Methyl (±)-7-[[[(6-amino-3-pyridinyl)amino]carbonyl]-3-oxo-4-(2-phenylethyl)-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate

Following the procedure of Example 1(a), substituting methyl (±)-7-carboxy-3-oxo-2-(2-phenylethyl)-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate for methyl (±)-7-carboxy-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate, and substituting 2,5-diaminopyridine for 2,6-diaminopyridine, the title compound was prepared (0.42 g, 42%): ¹H NMR (250 MHz, CDCl₃/DMSO-d₆) δ 6.14-9.46 (m, 7H), 5.35-5.44 (br, 3H), 5.13 (m, 1H),
25 3.68 (s, 3H), 3.55-3.83 (m, 3H), 2.94 (dd, J=17, 8 Hz, 1H), 2.78 (t, 2H), 2.66 (dd, J=17, 5 Hz, 1H); MS (ES) m/e 474.2 (M+H)⁺.
30

- b) (±)-7-[[[(6-Amino-3-pyridinyl)amino]carbonyl]-3-oxo-4-(2-phenylethyl)-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

35 1.0 N LiOH (1.1 mL, 1.1 mmol) was added dropwise to a mixture of methyl (±)-7-[[[(6-amino-3-pyridinyl)amino]carbonyl]-3-oxo-4-(2-phenylethyl)-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (0.23 g, 0.48 mmol) in THF (12 mL)

- 28 -

and H₂O (17 mL) at RT. After 22.5 h, the dark reddish-brown reaction mixture was concentrated on the rotavap to remove excess THF. The resulting reddish solution was cooled in an ice-bath, and neutralized with 1.0 N AcOH (1.8 mL). The precipitate was collected by suction filtration, washed with water and ether, and air-dried to give the title compound (0.19 g, 86%) as a pinkish-purple solid: ¹H NMR (400 MHz, DMSO-d₆) δ 9.55 (s, 1H), 6.40-8.29 (m, 7H), 5.69 (s, 2H), 5.42 (d, J=17 Hz, 1H), 5.09 (br, 1H), 3.95 (dd, J=17, 1H), 3.59-3.65 (m, 2H), 2.51-2.82 (m, 4H); MS (ES) m/e 460.2 (M+H)⁺. Anal. Calcd for C₂₅H₂₅N₅O₄ · 9/4 H₂O: C, 60.05; H, 5.98; N, 14.01. Found: C, 59.79; H, 5.92; N, 13.64.

10

Example 4

Preparation of (±)-1-Acetyl-7-[[6-amino-2-pyridinyl]aminocarbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

15

a) Methyl (±)-1-acetyl-7-carboxy-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate

Methyl (±)-7-carboxy-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (2.0 g, 8.6 mmol), acetic anhydride (25 mL) and acetic acid (1 mL) were heated to reflux. After 48 h, the reaction was concentrated, and the residue was diluted with H₂O (20 mL) and MeOH (15 mL). The resulting solution was concentrated to give the title compound (1.8 g, 78%): ¹H NMR (400 MHz, DMSO-d₆) δ 1.7-1.8 (s, 3H), 2.2-2.6 (m, 2H), 2.9-3.0 (s, 3H), 3.5-3.6 (s, 3H), 4.2-4.3 (d, 1H), 4.7-4.8 (d, 1H), 5.8-5.9 (t, 1H), 7.5-7.6 (d, 1H), 7.9-8.0 (dd, 1H), 8.1-8.2 (d, 1H); MS (ES) m/e 335.0 (M+H)⁺.

25

b) Methyl (±)-1-acetyl-7-[[6-amino-2-pyridinyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate

HOBT · H₂O (0.54 g, 3.6 mmol) and EDC (0.686 g, 3.6 mmol) were added to a solution of methyl (±)-1-acetyl-7-carboxy-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (1.0 g, 3 mmol) in dry DMF (15 mL). After 1 h the reaction solution was added dropwise to a solution of 2,6-diaminopyridine (0.326 g, 3.0 mmol) in dry DMF (20 mL). The reaction solution was stirred for 18 h, then was concentrated. Chromatography (silica gel, 9:1 CH₂Cl₂/MeOH) gave the title compound (0.9 g, 70%): ¹H NMR (400 MHz, DMSO-d₆) δ 1.80 (s, 3H), 2.3-2.7 (m, 2H), 3.0-3.1 (s, 3H), 3.5-3.6 (s, 3H), 4.1-4.2 (d, 1H), 4.7-4.8 (d, 1H),

35

- 29 -

5.7-5.9 (m, 3H), 6.2-6.3 (d, 1H), 7.3-7.5 (m, 3H), 8.0-8.1 (d, 1H), 8.2-8.3 (s, 1H);
MS (ES) m/e 426.0 (M+H)⁺.

- c) (±)-1-Acetyl-7-[[[(6-amino-2-pyridinyl)amino]carbonyl]-4-methyl-3-oxo-
5 2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid
1 N NaOH (1.0 mL, 1.0 mmol) was added to a cold solution of methyl (±)-
1-acetyl-7-[[[(6-amino-2-pyridinyl)amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-
tetrahydro-1H-1,4-benzodiazepine-2-acetate (0.23 g, 0.54 mmol), MeOH (2 mL)
and H₂O (1 mL). The solution was stirred at room temperature for 18 h then was
10 concentrated. Chromatography (ODS, 10% CH₃CN/H₂O-0.1% TFA) gave the title
compound (0.215 g, 90%): ¹H NMR (400 MHz, DMSO-d₆) δ 1.7-1.9 (s, 3H), 2.0-
2.6 (m, 2H), 3.0-3.1 (s, 3H), 4.1-4.2 (d, 1H), 4.7-4.8 (d, 1H), 5.7-5.9 (m, 1H), 6.4-
6.5 (d, 1H), 7.0-7.1 (d, 1H), 7.5-7.6 (d, 1H), 7.6-7.8 (t, 1H), 8.0-8.1 (d, 1H), 8.2-8.3
(s, 1H); MS (ES) m/e 412.2 (M+H)⁺. Anal. Calcd for C₂₀H₂₁N₅O₅ · 2.75
15 CF₃CO₂H: C, 42.25; H, 3.30; N, 9.66. Found: C, 42.01; H, 3.25; N, 9.85.

Example 5

- Preparation of (±)-2-methyl-3-oxo-8-[[2-(pyridinyl)carbonyl]amino]-2,3,4,5-
20 tetrahydro-1H-2-benzazepine-4-acetic acid

a) Methyl (±)-8-amino-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-
acetate

- A mixture of methyl (±)-8-[(benzyloxycarbonyl)amino]-2-methyl-3-oxo-
25 2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetate (1.5 g), Pearlman's catalyst (0.30 g),
and warm AcOH (300 mL) was stirred under H₂ (balloon pressure). After 90 min,
the mixture was filtered to remove the catalyst, and the filtrate was concentrated to
give the title compound as a viscous oil (1 g, 100%): ¹H NMR (CDCl₃) δ 7.00 (br
s, 2H), 6.88 (d, J=8.1 Hz, 1H), 6.57 (dd, J=8.1, 2.3 Hz, 1H), 6.43 (d, J=2.3 Hz, 1H),
30 5.25 (d, J=16.3 Hz, 1H), 3.78 (m, 1H), 3.70 (s, 3H), 3.68 (d, J=16.3 Hz, 1H), 3.03
(s, 3H), 3.00-2.80 (m, 3H), 2.40 (dd, J=16.7, 5.3 Hz, 1H).

b) Methyl (±)-2-methyl-3-oxo-8-[[2-(pyridinyl)carbonyl]amino]-2,3,4,5-
tetrahydro-1H-2-benzazepine-4-acetate

- 35 To a solution stirred under argon at room temperature of methyl (±)-8-
carboxy-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetate (0.30 g, 1
mmole), picolinic acid (1.2 mmole), HOBT · H₂O (0.17 g, 1.2 mmole),

- 30 -

diisopropylethylamine (0.53 g, 4 mmole), and DMF (5 mL) was added EDC (0.24 g, 1.2 mmole). The resulting mixture was stirred for 18 h, then was concentrated to dryness, and the residue was partitioned between EtOAc and H₂O. The organic phase was washed with H₂O and brine, dried (MgSO₄), and concentrated.

- 5 Purification by chromatography (silica gel, 1-5% MeOH/CH₂Cl₂) gave the title compound as an amorphous solid (26%): ¹H NMR (CDCl₃) δ 10.02 (s, 1H), 8.62 (d, J=4.1 Hz, 1H), 8.29 (d, J=7.8 Hz, 1H), 7.93 (m, 1H), 7.79 (d, J=2.1 Hz, 1H), 7.50 (m, 2H), 7.11 (d, J=8.3 Hz, 1H), 5.33 (d, J=16.6 Hz, 1H), 3.88 (d, J=16.6 Hz, 1H), 3.85 (m, 1H), 3.72 (s, 3H), 3.06 (s, 3H), 3.00-2.85 (m, 3H), 2.43 (dd, J=16.7, 10 5.4 Hz, 1H).

c) (±)-2-Methyl-3-oxo-8-[[2-(pyridinyl)carbonyl]amino]-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid

- A solution of methyl (±)-8-amino-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetate (0.054 g), LiOH · H₂O (0.008 g), THF (2 mL), and H₂O (2 15 mL) was stirred at RT. After 18 h, the reaction was concentrated to dryness and the residue was dissolved in H₂O. The solution was brought to pH 5 with 3N HCl, and the resulting precipitate was collected by filtration. Drying under vacuum gave the title compound as a colorless solid (0.04 g, 77%): ¹H NMR (DMSO-d₆) δ 8.77 (d, 20 J=4 Hz, 1H), 8.24 (d, J=8 Hz, 1H), 8.10 (m, 1H), 7.88 (d, J=2 Hz, 1H), 7.80-7.70 (m, 2H), 7.15 (d, J=8 Hz, 1H), 5.35 (d, J=16 Hz, 1H), 4.00 (d, J=16 Hz, 1H), 3.82 (m 1H), 2.97 (s, 3H), 2.78 (m, 2H), 2.55 (m, 1H), 2.38 (dd, J=16, 5 Hz, 2H); MS (ES) m/e 354 (M+H)⁺. Anal. Calcd for C₁₉H₁₉N₃O₄ · 0.125 H₂O: C, 64.17; H, 5.46; N, 11.82. Found: C, 63.99; H, 5.39; N, 11.77.

25

Example 6

Preparation of (±)-8-[(benzyloxycarbonyl)amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid

30

- a) Methyl (±)-8-[(benzyloxycarbonyl)amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetate

- To a solution stirred under argon at room temperature of methyl (±)-8-carboxy-2-methyl-3-oxo-2,3,4,5-tetrahydro-2-benzazepine-4-acetate (0.30 g, 1 35 mmol) and triethylamine (0.16 mL, 1.2 mmol) in toluene was added diphenylphosphorylazide (0.31 g, 1.2 mmol). The resulting mixture was heated at 80°C for 1 h, then was allowed to cool, and benzyl alcohol (0.22 g, 2 mmol) was

- 31 -

added. The mixture was heated at 80°C for 4 h, then was concentrated to remove the toluene. The residue was dissolved in EtOAc, and the solution was washed sequentially with 5% aqueous NaHCO₃, 1N HCl, and brine. Drying (MgSO₄), concentration, and chromatography (silica gel, 1% MeOH/CH₂Cl₂) gave the title compound as a colorless solid (0.20 g, 47%): ¹H NMR (CDCl₃) δ 7.40-7.30 (m, 6H), 7.06 (dd, J=8.3, 2 Hz, 1H), 7.02 (d, J=8.3 Hz, 1H), 6.65 (br s, 1H), 5.28 (d, J=16.7 Hz, 1H), 5.20 (s, 2H), 3.80 (m, 1H), 3.79 (d, J=16.7 Hz, 1H), 3.70 (s, 3H), 3.03 (s, 3H), 3.00 (m, 2H), 2.88 (m, 1H), 2.41 (dd, J=16.8, 5.3, 1H).

10 b) (±)-8-[(Benzyloxycarbonyl)amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid

A solution of the compound of Example 6(a) (0.24 mmol), LiOH · H₂O (0.013 g, 0.31 mmole), THF (2 mL), and H₂O (2 mL) was stirred at RT for 18 h, then was concentrated to dryness. The residue was dissolved in H₂O, and the solution was brought to pH 4-5 with 3N HCl. The resulting precipitate was collected by filtration and dried to yield the title compound (83%): ¹H NMR (Acetone-d₆) δ 8.50 (br s, 1H), 7.45 (br s, 1H), 7.40-7.30 (m, 5H), 7.05 (d, J=8.3 Hz, 1H), 5.31 (d, J=16.4 Hz, 1H), 5.17 (s, 2H), 3.90 (d, J=16.5 Hz, 1H), 3.84 (m, 1H), 3.08 (dd, J=11.6, 4.9 Hz, 1H); MS (ES) m/e 383.2 (M+H)⁺. Anal. Calcd for C₂₁H₂₂N₂O₅ · 0.25 H₂O: C, 65.19; H, 5.86; N, 7.24. Found: C, 65.45; H, 5.90; N, 7.27.

Example 7

25 Preparation of (±)-7-[[[3-(aminoiminomethyl)phenyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

a) Methyl (±)-7-[[[3-[N-(benzyloxycarbonyl)aminoiminomethyl]phenyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

30 Following the procedure of Example 2(a), methyl (±)-7-carboxy-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate was converted to the corresponding acid chloride. This was dissolved in CH₂Cl₂ (10 mL) and added dropwise to a solution of 3-amino-(N-benzyloxycarbonyl)benzamidine (1.48 g, 5.46 mmol) and pyridine (0.72 g, 9.1 mmol) in CH₂Cl₂ (15 mL) at 0°C under argon. The mixture was stirred in RT for 2 h, then was partitioned between CH₂Cl₂, EtOAc and 5% NaHCO₃. The organic layer was separated and washed with 0.05 N HCl, which gave a precipitate. This was collected and recrystallized

- 32 -

from MeOH/CH₂Cl₂/EtOAc to afford the title compound (0.5 g, 51%): ¹H NMR (400 MHz, CDCl₃/DMSO-d₆) δ 6.64-10.63 (m, 12H), 5.55 (d, J=16 Hz, 1H), 5.38 (s, 2H), 5.19 (m, 1H), 3.86 (d, J=17 Hz, 1H), 3.61 (s, 3H), 3.02 (s, 3H), 2.93 (dd, J=17, 9 Hz, 1H), 2.66 (dd, J=17, 4 Hz, 1H).

5

b) Methyl (±)-7-[[[3-(aminoiminomethyl)phenyl]amino] carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

A mixture containing the compound of Example 7(a), 10% Pd/C (0.19 g), and MeOH (30 mL) was hydrogenated at 45 psi in a Parr apparatus. After 2 h, glacial AcOH (5 mL) and more 10% Pd/C (0.15 g) were added, and hydrogenation at 45 psi was continued for an additional 70 min. The reaction mixture was filtered through a bed of Celite® and concentrated to give the title compound (0.21 g, 81%) as a white solid: ¹H NMR (250 MHz, DMSO-d₆) δ 6.55-10.19 (m, 10H), 5.55 (d, J=16 Hz, 1H), 5.18 (m, 1H), 3.90 (d, J=17 Hz, 1H), 3.62 (s, 3H), 2.97 (s, 3H), 2.86 (dd, J=17, 9 Hz, 1H), 2.67 (dd, J=17, 5 Hz, 1H); MS (ES) m/e 410.2 (M+H)⁺.

15

c) (±)-7-[[[3-(Aminoiminomethyl)phenyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

Following the procedure of Example 2(b), the compound of Example 7(b) was saponified and purified to give the title compound: ¹H NMR (250 MHz, DMSO-d₆) δ 6.51-9.30 (m, 10H), 5.54 (d, J=17 Hz, 1H), 5.14 (m, 1H), 3.88 (d, J=17 Hz, 1H), 2.98 (s, 3H), 2.82 (dd, J=17, 9 Hz, 1H), 2.57 (dd, J=17, 5 Hz, 1H); MS (ES) m/e 396.2 (M+H)⁺. Anal. Calcd for C₂₀H₂₁N₅O₄ · 3/2 CF₃CO₂H: C, 48.77; H, 4.00; N, 12.36. Found: C, 48.77; H, 4.38; N, 12.52.

25

Example 8

Preparation of (±)-7-[[[3-(aminocarbonyl)phenyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

The title compound was isolated as a minor product from the reaction described in Example 7(c): ¹H NMR (250 MHz, DMSO-d₆) δ 9.91 (s, 1H), 6.42-8.25 (m, 9H), 5.53 (d, J=17 Hz, 1H), 5.12 (m, 1H), 3.87 (d, J=17 Hz, 1H), 2.98 (s, 3H), 2.82 (dd, J=17, 9 Hz, 1H), 2.57 (dd, J=17, 5 Hz, 1H); MS (ES) m/e 397.0 (M+H)⁺. Anal. Calcd for C₂₀H₂₀N₅O₅ · 1/3 CF₃CO₂H · 2/3 H₂O: C, 55.60; H, 4.89; N, 12.55. Found: C, 55.38; H, 5.05; N, 12.42.

35

Example 9Preparation of (±)-4-methyl-3-oxo-7-[1-(4-phenylpiperazinyl)carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

5

a) Methyl (±)-4-methyl-3-oxo-7-[1-(4-phenylpiperazinyl)carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate

EDC (137.6 mg, 0.72 mmol) was added to a solution of methyl (±)-7-carboxy-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate
10 (175.4 mg, 0.6 mmol), 4-phenylpiperazine (0.11 mL, 0.72 mmol), HOBT·H₂O (97.3 mg, 0.72 mmol), and diisopropylethylamine (0.21 mL, 1.2 mmol) in anhydrous DMF (3 mL) at RT. After 18 h, the reaction was concentrated on the rotavap (high vacuum), and the residue was diluted with H₂O. CHCl₃ extraction drying (Na₂SO₄), concentration, and reconcentration from xylenes left a yellow oil,
15 which solidified on treatment with EtOAc followed by concentration. Chromatography (silica gel, 5% MeOH/CHCl₃) gave an oil which was treated with EtOAc to afford a solid. Trituration with cold Et₂O (2 x 20 mL) gave the title compound as an off-white solid (238.8 mg, 91%): TLC R_f (5% MeOH/CHCl₃) 0.33; ¹H NMR (250 MHz, CDCl₃) δ 7.07-7.36 (m, 4H), 6.82-7.00 (m, 3H), 6.49-
20 6.59 (m, 1H), 5.45 (d, J=16.5 Hz, 1H), 5.19 (app t, 1H), 3.55-3.95 (m, 5H), 3.75 (s, 3H), 3.15-3.35 (m, 4H), 3.09 (s, 3H), 3.00 (dd, J=16.0, 6.6 Hz, 1H), 2.67 (dd, J=16.0, 6.5 Hz, 1H); MS (ES) m/e 437.2 (M+H)⁺, 275.0.

b) (±)-4-Methyl-3-oxo-7-[1-(4-phenylpiperazinyl)carbonyl]-2,3,4,5-tetrahydro-1H-
25 1,4-benzodiazepine-2-acetic acid

A mixture of methyl (±)-4-methyl-3-oxo-7-[1-(4-phenylpiperazinyl)carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (238.8 mg, 0.55 mmol), 1.0 N NaOH (1.7 mL, 1.7 mmol), and MeOH (5.5 mL) was stirred at 35°C overnight. The resulting homogeneous solution was concentrated to
30 dryness, and the residue was dissolved in 1:1 H₂O/CH₃CN. The solution was cooled to 0°C and acidified with TFA, then was concentrated to a pinkish-orange oil. Chromatography (ODS, 25% CH₃CN/H₂O-0.1% TFA), concentration, and lyophilization gave the title compound (213.1 mg, 69%) as an off-white powder: HPLC k' 2.6 (PRP-1®, 25% CH₃CN/H₂O-0.1% TFA); ¹H NMR (400 MHz, CD₃OD) δ 7.29-7.40 (m, 2H), 7.12-7.22 (m, 4H), 7.04 (t, J=7.4 Hz, 1H), 6.63 (d, J=9.1 Hz, 1H), 5.58 (d, J=16.5 Hz, 1H), 5.20 (dd, J=9.0, 5.1 Hz, 1H), 3.78-3.95 (m, 5H), 3.23-3.39 (m, 4 H, partially obscured by residual solvent signal), 3.04 (s, 3H),

- 34 -

2.94 (dd, J=16.8, 9.0 Hz, 1H), 2.65 (dd, J=16.8, 5.1 Hz, 1H); MS (ES) m/e 423.2 (M+H)⁺, 261.0. Anal. Calcd for C₂₃H₂₆N₄O₄ · 1.25 CF₃CO₂H: C, 54.20; H, 4.86; N, 9.92. Found: C, 54.42; H, 4.98; N, 10.00.

5

Example 10Preparation of (±)-4-methyl-3-oxo-7-[(1-piperazinyl)carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

- 10 a) Methyl (±)-4-methyl-3-oxo-7-[1-[4-(tert-butoxycarbonyl)piperazinyl]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate

EDC (137.6 mg, 0.72 mmol) was added to a solution of methyl (±)-7-carboxy-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (175.4 mg, 0.6 mmol), 4-(tert-butoxycarbonyl)piperazine (0.21 g, 1.2 mmol),
15 HOBT·H₂O (97.3 mg, 0.72 mmol), and diisopropylethylamine (0.21 mL, 1.2 mmol) in anhydrous DMF (3 mL) at RT. After 17 h, the reaction was concentrated on the rotavap (high vacuum), and the residue was partitioned between 10% Na₂CO₃ and CH₂Cl₂. The layers were separated and the aqueous layer was extracted with CH₂Cl₂. Drying (Na₂SO₄) and concentration left a residue which
20 was reconcentrated from xylenes to remove DMF. Chromatography (silica gel, 10% MeOH in 1:1 EtOAc/CHCl₃) gave the title compound as a colorless foam (266 mg, 96%): TLC R_f 0.54 (10% MeOH in 1:1 EtOAc/CHCl₃); ¹H NMR (250 MHz, CDCl₃) δ 7.09-7.16 (m, 2H), 6.53 (d, J=8.9 Hz, 1H), 5.44 (d, J=16.5 Hz, 1H), 5.07 (t, J=6.6 Hz, 1H), 3.75 (s, 3H), 3.73 (d, J=16.5 Hz, 1H, partially
25 obscured by the δ 3.75 resonance), 3.32-3.68 (m, 8H), 3.08 (s, 3H), 2.99 (dd, J=16.0, 6.7 Hz, 1H), 2.67 (dd, J=16.0, 6.4 Hz, 1H), 1.47 (s, 9H); IR (CHCl₃) 3240-3560 (br), 1729, 1685 (shoulder), 1665, 1610, 1455, 1437, 1420, 1410, 1367, 1288, 1263, 1245, 1168, 1123 cm⁻¹; MS (ES) m/e 461.2 (M+H)⁺, 405.2 (M+H-C₄H₈)⁺.

- 30 b) (±)-4-Methyl-3-oxo-7-[(1-piperazinyl)carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

TFA (2.9 mL) was added all at once to a solution of methyl (±)-4-methyl-3-oxo-7-[1-[4-(tert-butoxycarbonyl)piperazinyl]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (266.2 mg, 0.58 mmol) in anhydrous CH₂Cl₂ (2.9 mL) at
35 0°C, and the reaction was warmed to RT. After 2 h, the reaction was concentrated and the residue was reconcentrated several times from 1:1 toluene/MeOH to remove residual TFA. The resulting yellow oil was dissolved in MeOH (5.8 mL)

- 35 -

and cooled to 0°C. 1.0 N NaOH (2.9 mL, 2.9 mmol) was added, and the bright yellow solution was stirred at RT. After 20 h, the reaction was concentrated and the residue was dissolved in H₂O (5 mL). The solution was acidified with TFA and concentrated to leave a light yellow oil. Chromatography (ODS, 5% CH₃CN/H₂O-0.1% TFA), concentration, and lyophilization gave the title compound as a light yellow powder (206.1 mg, 66%): HPLC k' 2.2 (PRP-1®, 7% CH₃CN/H₂O-0.1% TFA); ¹H NMR (400 MHz, CD₃OD) δ 7.15-7.23 (m, 2H), 6.59-6.65 (m, 1H), 5.58 (d, J=16.5 Hz, 1H), 5.20 (dd, J=9.0, 5.1 Hz, 1H), 3.81-3.95 (m, 5H), 3.22-3.30 (m, 4H), 3.04 (s, 3H), 2.94 (dd, J=16.7, 9.0 Hz, 1H), 2.65 (dd, J=16.7, 5.1 Hz, 1H); MS (ES) m/e 347.2 (M+H)⁺, 261.0. Anal. Calcd for C₁₇H₂₂N₄O₄ · 1.5 CF₃CO₂H · H₂O: C, 44.86; H, 4.80; N, 10.46. Found: C, 44.82; H, 4.82; N, 10.36.

Example 11

15 Preparation of (±)-4-methyl-3-oxo-7-[1-[4-(1-piperidiny)l]piperidiny]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

a) Methyl (±)-4-methyl-3-oxo-7-[1-[4-(1-piperidiny)l]piperidiny]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate

20 EDC (138 mg, 0.72 mmol) was added to a solution of methyl (±)-7-carboxy-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (175 mg, 0.6 mmol), 4-(1-piperidiny)l]piperidine (121.2 mg, 0.72 mmol), HOBT·H₂O (97 mg, 0.72 mmol), and diisopropylethylamine (0.21 mL, 1.2 mmol) in anhydrous DMF (3 mL) at RT. After 21 h, the reaction was concentrated on the rotavap (high vacuum), and the residue was taken up in CH₂Cl₂ and washed with H₂O. Drying (Na₂SO₄) and concentration left the crude product as an off-white solid, which was reconstituted from xylenes to remove residual DMF. The aqueous washings were combined and extracted with CHCl₃. The combined CHCl₃ layers were dried (Na₂SO₄), concentrated, and reconstituted from xylenes to provide additional crude product. The crude products were combined and chromatographed (silica gel, 20% MeOH/CHCl₃) to give the title compound as a colorless solid (205.2 mg, 77%): TLC R_f 0.39 (20% MeOH/CHCl₃); ¹H NMR (250 MHz, CDCl₃) δ 7.05-7.15 (m, 2H), 6.52 (d, J=8.8 Hz, 1H), 5.44 (d, J=16.4 Hz, 1H), 5.00-5.12 (m, 1H), 4.10-4.70 (m, 2H), 4.40 (d, J=5.1 Hz, 1H), 3.75 (s, 3H), 3.73 (d, J=16.4 Hz, 1H, partially obscured by the δ 3.75 resonance), 3.08 (s, 3H), 2.40-3.20 (m, 9H), 1.35-2.10 (m, 10H); IR (CHCl₃) 3400, 2930, 1731, 1669, 1613, 1439, 1282 cm⁻¹; MS (ES) m/e 443.2 (M+H)⁺.

b) (±)-4-Methyl-3-oxo-7-[1-[4-(1-piperidiny)l]piperidiny]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

- A mixture of methyl (±)-4-methyl-3-oxo-7-[1-[4-(1-piperidiny)l]piperidiny]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (205.2 mg, 0.48 mmol), 1.0 N NaOH (1.4 mL, 1.4 mmol), and MeOH (4.8 mL) was stirred at 35-40°C for 19.5 h, then was concentrated to dryness. The residue was dissolved in H₂O (2-3 mL), cooled to 0°C, and acidified with TFA. Concentration left a residue which was purified by chromatography (ODS, step gradient, 12% CH₃CN/H₂O-0.1% TFA, 15% CH₃CN/H₂O-0.1% TFA). Concentration and lyophilization gave the title compound as a colorless powder (229.1 mg, 78%): HPLC k' 2.9 (PRP-1®, 12% CH₃CN/H₂O-0.1% TFA); ¹H NMR (400 MHz, CD₃OD) δ 7.11-7.18 (m, 2H), 6.61 (d, J=9.0 Hz, 1H), 5.58 (d, J=16.6 Hz, 1H), 5.20 (dd, J=9.0, 5.1 Hz, 1H), 4.25-4.68 (m, 2H), 3.88 (d, J=16.6 Hz, 1H), 3.41-3.56 (m, 3H), 3.04 (s, 3H), 2.88-3.16 (m, 4H), 2.94 (dd, J=16.7, 9.0 Hz, 1H), 2.65 (dd, J=16.7, 5.1 Hz, 1H), 1.43-2.20 (m, 10H); MS (ES) m/e 429.2 (M+H)⁺. Anal. Calcd for C₂₃H₃₂N₄O₄ · 1.5 CF₃CO₂H · 0.5 H₂O: C, 51.31; H, 5.71; N, 9.21. Found: C, 51.51; H, 5.90; N, 9.29.

20

Example 12

Preparation of (±)-4-methyl-3-oxo-7-[1-[4-(2-pyridiny)l]piperaziny]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

- a) Methyl (±)-4-methyl-3-oxo-7-[1-[4-(2-pyridiny)l]piperaziny]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate
- EDC (137.6 mg, 0.72 mmol) was added to a solution of methyl (±)-7-carboxy-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (175.4 mg, 0.6 mmol), 4-(2-pyridiny)l]piperazine (0.11 mL, 0.72 mmol), HOBT·H₂O (97.3 mg, 0.72 mmol), and diisopropylethylamine (0.21 mL, 1.2 mmol) in anhydrous DMF (3 mL) at RT. After 18.5 h, the mixture was concentrated on the rotavap (high vacuum), and the residue was partitioned between 10% aqueous Na₂CO₃ and CHCl₃. The layers were separated and the aqueous layer was extracted with CHCl₃. The combined CHCl₃ layers were washed with brine, dried (Na₂SO₄), and concentrated. The residue was concentrated from 1:1:1 CHCl₃/MeOH/xylene to remove DMF, then was chromatographed (silica gel, 5% MeOH/CHCl₃). The impure oily product obtained

- 37 -

in this way was dissolved in EtOAc. Crystallization began almost immediately, and the mixture was cooled in ice. The solid was collected by suction filtration and was washed with EtOAc. Drying in high vacuum gave the title compound as a nearly colorless solid (239.6 mg, 91%): mp >250°C; TLC R_f 0.39 (5%

5 MeOH/CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 8.18-8.25 (m, 1H), 7.49-7.59 (m, 1H), 7.15-7.23 (m, 2H), 6.65-6.75 (m, 2H), 6.54 (d, J=8.0 Hz, 1H), 5.45 (d, J=16.5 Hz, 1H), 5.03-5.12 (m, 1H), 4.40 (d, J=5.0 Hz, 1H), 3.68-3.85 (m, 5H), 3.75 (s, 3H), 3.51-3.68 (m, 4H), 3.09 (s, 3H), 3.00 (dd, J=16.0, 6.6 Hz, 1H), 2.68 (dd, J=16.0, 6.5 Hz, 1H); MS (ES) m/e 438.2 (M+H)⁺, 275.0.

10

b) (±)-4-Methyl-3-oxo-7-[1-[4-(2-pyridinyl)piperazinyl]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

1.0 N NaOH (0.65 mL, 0.65 mmol) was added dropwise to a suspension of methyl (±)-4-methyl-3-oxo-7-[1-[4-(2-pyridinyl)piperazinyl]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (235 mg, 0.54 mmol) in MeOH (5.4 mL) at RT, and the resulting mixture was stirred at 50°C. After 3.5 h, more 1.0 N NaOH (0.65 mL, 0.65 mmol) was added, and warming at 50°C was continued for 0.5 h. The resulting homogeneous solution was stirred at RT for 18.5 h, then was concentrated. The residue was dissolved in H₂O (5 mL), and the solution was
20 acidified with TFA. Concentration left a foamy oil which was purified by chromatography (ODS, 12% CH₃CN/H₂O-0.1% TFA). Concentration and lyophilization gave the title compound (280 mg, 87%) as a colorless powder: HPLC k' 3.7 (PRP-1®, 12% CH₃CN/H₂O-0.1% TFA); ¹H NMR (400 MHz, CD₃OD) δ 7.96-8.10 (m, 2H), 7.37 (d, J=9.1 Hz, 1H), 7.28-7.38 (m, 2H), 7.02 (app
25 t, 1H), 6.64 (d, J=9.1 Hz, 1H), 5.58 (d, J=16.5 Hz, 1H), 5.21 (dd, J=9.0, 5.1 Hz, 1H), 3.68-3.97 (m, 9H), 3.04 (s, 3H), 2.94 (dd, J=16.8, 9.0 Hz, 1H), 2.66 (dd, J=16.8, 5.1 Hz, 1H); MS (ES) m/e 424.2 (M+H)⁺. Anal. Calcd for C₂₂H₂₅N₅O₄ · 1.5 CF₃CO₂H: C, 50.51; H, 4.49; N, 11.78. Found: C, 50.42; H, 4.78; N, 11.77.

30

Example 13

Preparation of (±)-4-methyl-3-oxo-7-[1-[4-(phenylmethyl)piperazinyl]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

- 38 -

a) Methyl (±)-4-methyl-3-oxo-7-[1-[4-(phenylmethyl)piperazinyl]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate

Methyl (±)-7-carboxy-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (292.3 mg, 1.0 mmol) was refluxed with thionyl chloride (10 mL) for 20 min, and the yellow solution was concentrated. The residue was concentrated from dry toluene (5 mL), then was taken up in dry CH₂Cl₂ (10 mL) and cooled to 0°C under argon. 1-benzylpiperazine (0.70 mL, 4 mmol) was added all at once, and after 5 min, the light yellow solution was warmed to RT. The reaction was stirred for 0.5 h, then was washed sequentially with 1.0 N NaOH and H₂O. Drying (MgSO₄), concentration, and chromatography (silica gel, 10% MeOH in 1:1 EtOAc/CHCl₃) gave the title compound as an off-white foam (344.4 mg, 76%): TLC R_f 0.47 (10% MeOH in 1:1 EtOAc/CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 7.22-7.40 (m, 5H), 7.08-7.20 (m, 2H), 6.50 (d, J=8.8 Hz, 1H), 5.43 (d, J=16.4 Hz, 1H), 5.02-5.10 (m 1H), 4.37 (br d, J=5.1 Hz, 1H), 3.45-3.85 (m, 7H), 3.74 (s, 3H), 3.07 (s, 3H), 2.98 (dd, J=16.1, 6.7 Hz, 1H), 2.66 (dd, J=16.1, 6.4 Hz, 1H); FTIR (CCl₄) 1740, 1672, 1630, 1611, 1456, 1437, 1296 cm⁻¹; MS (ES) m/e 451.2 (M+H)⁺.

b) (±)-4-Methyl-3-oxo-7-[1-[4-(phenylmethyl)piperazinyl]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

1.0 N LiOH (0.91 mL, 0.91 mmol) was added to a solution of methyl (±)-4-methyl-3-oxo-7-[1-[4-(phenylmethyl)piperazinyl] carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (344.4 mg, 0.76 mmol) in THF (3.8 mL) and H₂O (2.9 mL) at 0°C. The resulting bright yellow solution was stirred at RT for 15 h, then was acidified with TFA (0.23 mL) and concentrated to dryness. Chromatography (ODS, 20% CH₃CN/H₂O-0.1% TFA), concentration, and lyophilization gave the title compound as a colorless powder (349.9 mg, 77%): HPLC k' 1.6 (PRP-1®, 20% CH₃CN/H₂O-0.1% TFA); ¹H NMR (400 MHz, CD₃OD) δ 7.43-7.57 (m, 5H), 7.14-7.22 (m, 2H), 6.62 (d, J=9.0 Hz, 1H), 5.57 (d, J=16.6 Hz, 1H), 5.20 (dd, J=9.0, 5.1 Hz, 1H), 4.37 (s, 2H), 3.88 (d, J=16.6 Hz, 1H), 3.45-4.30 (m, 2H), 3.15-3.45 (m, 6 H, partially obscured by residual solvent signal), 3.03 (s, 3H), 2.93 (dd, J=16.9, 9.0 Hz, 1H), 2.65 (dd, J=16.9, 5.1 Hz, 1H); MS (ES) m/e 437.2 (M+H)⁺. Anal. Calcd for C₂₄H₂₈N₄O₄ · 1.25 CF₃CO₂H: C, 53.31; H, 5.28; N, 9.38. Found: C, 53.30; H, 5.42; N, 9.35.

35

Example 14Preparation of (±)-4-methyl-3-oxo-7-[1-[4-(2-pyrimidinyl)piperazinyl]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

5

a) Methyl (±)-4-methyl-3-oxo-7-[1-[4-(2-pyrimidinyl)piperazinyl]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate

A mixture of methyl (±)-7-carboxy-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (584 mg, 2.0 mmol), 1-(2-pyrimidinyl)piperazine dihydrochloride (522 mg, 2.2 mmol), HOBT·H₂O (270 mg, 2 mmol), triethylamine (1.0 mL, 7.2 mmol), and EDC (383 mg, 2 mmol) in anhydrous DMF (40 mL) was stirred at RT overnight. The reaction was concentrated in vacuum, and the resulting residue was diluted with 5% K₂CO₃. CH₂Cl₂ extraction, drying (MgSO₄), and concentration gave the title compound (0.76 g, 86%): MS (ES) m/e 483 (M+H)⁺.

15

b) (±)-4-Methyl-3-oxo-7-[1-[4-(2-pyrimidinyl)piperazinyl]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

Methyl (±)-4-methyl-3-oxo-7-[1-[4-(2-pyrimidinyl)piperazinyl]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (0.7 g, 1.6 mmol) was suspended in MeOH (10 mL) and THF (5 mL), and 1.0 N NaOH (6 mL) were added. The reaction was stirred at RT for 2 d, then was concentrated in vacuum. The residue was diluted with H₂O, and the pH was adjusted to 5 to 6 with 1.5 N HCl. Lyophilization gave the title compound (0.186 g, 27%): MS (ES) m/e 425 (M+H)⁺. Anal. Calcd for C₂₁H₂₄N₆O₄ · 2.4 H₂O: C, 53.93; H, 6.21; N, 17.97. Found: C, 54.25; H, 5.96; N, 17.53.

20

25

Example 15

30 Preparation of 8-[[2S-amino-3-phenylpropanoyl]amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4RS-acetic acid

a) Methyl-8-[[2S-(tert-butoxycarbonyl)amino-3-phenylpropanoyl]amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4RS-acetate

35

Methyl (±)-8-amino-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetate was coupled with Boc-L-phenylalanine according to the procedure of Example 5(b). Purification by chromatography (silica gel, 1%-5%

- 40 -

CH₃OH/CH₂Cl₂) gave the title compound as a light yellow foam (99%): ¹H NMR (CDCl₃) δ 7.80 (m, 1H), 7.30 (m, 5H), 7.02 (m, 2H), 5.25 (d, J=15.9 Hz, 1H), 5.10 (m, 1H), 4.45 (m, 1H), 3.80 (m, 1H), 3.78 (d, J=15.9 Hz, 1H), 3.71 (s, 3H), 3.14 (d, J=6.8 Hz, 1H), 3.03 (s, 3H), 2.97 (m, 2H), 2.88 (m, 1H), 2.40 (dd, J=16.8, 5.4 Hz, 1H), 1.43 (s, 9H).

b) 8-[[2S-Amino-3-phenylpropanoyl]amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4RS-acetic acid

Methyl-8-[[2S-(tert-butoxycarbonyl)amino-3-phenylpropanoyl]amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4(R,S)-acetate was deprotected according to the procedure of Example 20(b) to give the title compound as a light yellow foam (58%): MS (ES) m/e 396(M+H)⁺. Anal. Calcd for C₂₂H₂₅N₃O₄ · 1.57 C₂F₃HO₂: C, 52.56; H, 4.66; N, 7.73. Found: C, 52.95; H, 4.90; N, 6.92.

Example 16

Preparation of (±)-8-[[[(2-hydroxy-2-phenyl)ethyl]methylamino]carbonyl]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid

a) Methyl (±)-8-[[[(2-hydroxy-2-phenyl)ethyl]methylamino]carbonyl]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetate

EDC (230 mg, 1.2 mmole) was added to a stirred solution of methyl (±)-8-carboxy-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetate (1.0 mmol), (±)-α-(methylaminomethyl)benzyl alcohol (1.2 mmol), HOBT·H₂O (162 mg, 1.2 mmol), and diisopropylethylamine (0.70 mL, 4.0 mmol) in anhydrous DMF (5 mL) at RT. After 19 hr, the reaction was concentrated on the rotavap (high vacuum), and the residue was partitioned between H₂O and EtOAc. The layers were separated and the organic layer was washed with H₂O. Drying (MgSO₄), concentration, and chromatography (silica gel, 1%-3% CH₃OH/CH₂Cl₂) gave the title compound as an off-white foam (62%): ¹H NMR (CDCl₃) (mixture of diastereomers) δ 8.62 (d, J=5 Hz), 8.29 (d, J=7.8 Hz), 7.93 (m), 7.81 (m), 7.62 (s), 7.45 (m), 7.18 (d, J=5 Hz), 7.11 (d, J=8.3 Hz), 7.00 (d, J=8.2 Hz), 5.42 (br s), 5.33 (d, J=16), 5.25 (d, J=16 Hz), 3.88 (d, J=16.5 Hz), 3.85 (m), 3.79 (d, J=16.3 Hz), 3.70 (s), 3.69 (s), 3.05 (s), 3.01 (s), 3.00 (m), 2.43 (m), 2.18 (s).

- 41 -

b) (±)-8-[[[(2-Hydroxy-2-phenyl)ethyl]methylamino]carbonyl]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid

- Methyl (±)-8-[[[(2-hydroxy-2-phenyl)ethyl]methylamino]carbonyl]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetate was saponified according to the procedure of Example 6(b) to give the title compound as a white solid (80%): ¹H NMR (DMSO-d₆) δ 8.12 (m, 1H), 7.80 (m, 2H), 7.53 (m, 2H), 7.41 (m, 2H), 5.09 (m, 2H), 4.80 (m, 2H), 3.97 (m, 1H), 3.40 (m, 5H), 3.18 (s, 3H), 3.08 (m, 1H), 2.87 (m, 1H), 2.64 (m, 1H), 2.36 (m, 1H); MS (ES) m/e 411 (M+H)⁺. Anal. Calcd for C₂₃H₂₆N₂O₅ · 1.75 H₂O: C, 62.50; H, 6.73; N, 6.34. Found: C, 62.60; H, 6.63; N, 6.19.

Example 17

Preparation of (±)-7-[[[N-(2-hydroxyethyl)-N-methyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

- a) Methyl (±)-7-[[[N-(2-hydroxyethyl)-N-methyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate
- EDC (229.4 mg, 1.2 mmol) was added to a stirred solution of methyl (±)-7-carboxy-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (292.3 mg, 1.0 mmol), 2-(methylamino)ethanol (0.10 mL, 1.2 mmol), HOBT·H₂O (162.2 mg, 1.2 mmol), and diisopropylethylamine (0.35 mL, 2.0 mmol) in anhydrous DMF (5 mL) at RT. After 15.5 h, the reaction was concentrated on the rotavap (high vacuum), and the resulting yellow oil was dissolved in CHCl₃. The solution was washed with 10% Na₂CO₃, and the aqueous was back-extracted with CHCl₃. The combined organic layers were dried (Na₂SO₄), concentrated, and reconcentrated from xylenes to remove residual DMF. Chromatography (silica gel, 2:2:1 EtOAc/CHCl₃/MeOH) gave the title compound as a colorless oil (336 mg, 96%): TLC R_f 0.46 (2:2:1 EtOAc/CHCl₃/MeOH); ¹H NMR (250 MHz, CDCl₃) δ 7.15-7.25 (m, 2H), 6.51 (d, J=8.9 Hz, 1H), 5.44 (d, J=16.4 Hz, 1H), 5.08 (t, J=6.6 Hz, 1H), 3.52-3.98 (m, 5H), 3.75 (s, 3H), 3.10 (s, 3H), 3.08 (s, 3H), 3.00 (dd, J=16.0, 6.8 Hz, 1H), 2.67 (dd, J=16.0, 6.4 Hz, 1H); IR (CHCl₃) 3060-3540, 1729, 1659, 1612, 1482, 1437, 1402 cm⁻¹; MS (ES) m/e 350.0 (M+H)⁺.

- 42 -

b) (±)-7-[[[N-(2-Hydroxyethyl)-N-methyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

- 1.0 N NaOH (1.4 mL, 1.4 mmol) was added dropwise to a solution of methyl (±)-7-[[[N-(2-hydroxyethyl)-N-methyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (336.3 mg, 0.96 mmol) in MeOH (4.8 mL) at 0°C, and the resulting solution was stirred at RT. After 16 h, the reaction was concentrated, and the residue was purified by chromatography (ODS, 30% MeOH/H₂O). The fractions containing the product were combined, concentrated to dryness, and the residue was repurified by ODS chromatography (15% MeOH/H₂O). Concentration and lyophilization gave the title compound as a colorless powder (231 mg, 64%): HPLC k' 3.7 (PRP-1®, 10% CH₃CN/H₂O-0.1% TFA); ¹H NMR (400 MHz, CD₃OD) δ 7.11-7.20 (m, 2H), 6.59 (d, J=8.1 Hz, 1H), 5.59 (d, J=16.5 Hz, 1H), 5.15 (t, J=7.0 Hz, 1H), 3.85 (d, J=16.5 Hz, 1H), 3.45-3.85 (m, 4H), 3.10 (br s, 3H), 3.03 (s, 3H), 2.79 (dd, J=15.5, 7.5 Hz, 1H), 2.45 (dd, J=15.5, 6.4 Hz, 1H); MS (ES) m/e 358.0 (M+Na)⁺, 336.0 (M+H)⁺. Anal. Calcd for C₁₆H₂₀N₃O₅Na · H₂O: C, 51.20; H, 5.91; N, 11.19. Found: C, 51.04; H, 6.04; N, 11.14.

Example 18

20

Preparation of (±)-8-[[2-(2-pyridinylamino)acetyl]amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid

a) Methyl 2-(2-pyridinyl)acetate hydrochloride

- 25 A solution of 2-(2-pyridinyl)acetic acid (0.50 g, 3 mmol; prepared according to *J. Prakt. Chem.* 1961, 285, 118) in MeOH (20 mL) was saturated with HCl gas at 0°C, then was allowed to warm to RT. After stirring overnight, the reaction was concentrated under vacuum to afford the title compound (0.54 g, 100%) as a colorless solid: ¹H NMR (CDCl₃) δ 8.90 (br s, 1H), 7.89 (m, 2H), 6.89 (t, J=6.5 Hz, 1H), 6.79 (d, J=8 Hz, 1H), 4.21 (m, 2H), 3.81 (s, 3H).
- 30

b) Methyl 2-(N-tert-butoxycarbonyl-N-2-pyridinyl)aminoacetate

- Methyl 2-(2-pyridinyl)aminoacetate hydrochloride (1.0 g, 5 mmol) was partitioned between EtOAc and 10% K₂CO₃. The organic layer was washed with brine, dried (MgSO₄), and concentrated. The residue was dissolved in CH₂Cl₂ (10 mL), and di-tert-butyl dicarbonate (1.2 g, 5 mmol) was added. The reaction was stirred at RT under argon overnight, then was concentrated. Chromatography on
- 35

- 43 -

silica gel (1%-5% CH₃OH/CH₂Cl₂) gave the title compound as a pale yellow oil (0.89 g, 68%): ¹H NMR (CDCl₃) δ 8.31 (dd, J=4.8, 1.5 Hz, 1H), 7.84 (d, J=4.8 Hz, 1H), 7.67 (m, 1H), 7.00 (m, 1H), 4.73 (s, 2H), 3.75 (s, 3H), 1.52 (s, 9H).

5 c) 2-(N-tert-butoxycarbonyl-N-2-pyridinyl)aminoacetic acid

Methyl 2-(N-tert-butoxycarbonyl-N-2-pyridinyl)aminoacetate was saponified according to the procedure of Example 6(b) to give the title compound (74%) as a white solid.

10 d) Methyl (±)-8-[[2-(N-tert-butoxycarbonyl-N-2-pyridinyl)aminoacetyl]amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetate

Methyl (±)-8-amino-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetate was coupled with 2-(N-tert-butoxycarbonyl-N-2-pyridinyl)aminoacetic acid according to the procedure of Example 5(b). Purification by silica gel
15 chromatography (1%-3% CH₃OH/CH₂Cl₂) gave the title compound as a light yellow foam (56%): ¹H NMR (CDCl₃) δ 9.72 (s, 1H), 8.45 (d, J=4.2 Hz, 1H), 7.89 (t, J=4.2 Hz, 1H), 7.80 (d, J=8.3 Hz, 1H), 7.50 (s, 1H), 7.26 (m, 2H), 7.03 (d, J=8.4 Hz, 1H), 5.28 (d, J=16.4 Hz, 1H), 4.66 (s, 2H), 3.82 (d, J=16.4 Hz, 1H), 3.80 (m, 1H), 3.71 (s, 3H), 3.04 (s, 3H), 3.00 (m, 2H), 2.90 (m, 1H), 2.41 (dd, J=16.8, 5.2
20 Hz, 1H), 1.52 (s, 9H).

e) (±)-8-[[2-(2-pyridinylamino)acetyl]amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid

Methyl (±)-8-[[2-(N-tert-butoxycarbonyl-N-2-
25 pyridinyl)aminoacetyl]amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetate was deprotected according to the procedure of Example 20(b). Purification by chromatography (ODS, 10% CH₃CN/H₂O-0.1% TFA) followed by lyophilization gave the title compound as an off-white powder (23%): ¹H NMR (CD₃OD) δ 7.99 (m, 1H), 7.90 (d, J=6.4 Hz, 1H), 7.48 (d, J=2 Hz, 1H),
30 7.41 (dd, J=8.3, 2 Hz, 1H), 7.15 (d, J=9.2 Hz, 1H), 7.11 (d, J=8.4 Hz, 1H), 6.98 (m, 1H), 5.35 (d, J=16.4 Hz, 1H), 4.32 (s, 2H), 3.94 (d, J=16.4 Hz, 1H), 3.90 (m, 1H), 3.07 (dd, J=17, 4.3 Hz, 1H), 2.70 (m, 2H), 2.45 (dd, J=17, 5 Hz, 1H); MS (ES) m/e 383.2(M+H)⁺. Anal. Calcd for C₂₀H₂₂N₄O₄ · 1.5 C₂F₃HO₂ · 0.5 H₂O: C, 49.11; H, 4.39; N, 9.96. Found: C, 49.23; H, 4.27; N, 10.07.

35

Example 19Preparation of (±)-4-methyl-3-oxo-7-[[2-(phenylaminoethyl)amino]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

5

a) Methyl (±)-4-methyl-3-oxo-7-[[2-(phenylaminoethyl)amino]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate

Following the procedure of Example 14(a), except substituting N-phenylethylenediamine for the 1-(2-pyrimidinyl)piperazine dihydrochloride, the
10 title compound was prepared (78%): MS (ES) m/e 411 (M+H)⁺.

b) (±)-4-Methyl-3-oxo-7-[[2-(phenylaminoethyl)amino]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

Following the procedure of Example 14(b), methyl (±)-4-methyl-3-oxo-7-
15 [[2-(phenylaminoethyl)amino]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate was saponified to give the title compound (24%): MS (ES) m/e 397 (M+H)⁺. Anal. Calcd for C₂₁H₂₄N₄O₄ · 1.5 CF₃CO₂H · 1.5 H₂O: C, 48.49; H, 4.83; N, 9.42. Found: C, 48.41; H, 4.81; N, 9.40.

20

Example 20Preparation of (±)-8-[(2-aminoacetyl)amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1h-2-benzazepine-4-acetic acid

25 a) Methyl (±)-8-[(2-[tert-butoxycarbonyl]aminoacetyl)amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetate

Methyl (±)-8-amino-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetate was coupled with Boc-glycine according to the procedure of example 5(b). Purification by chromatography (silica gel, 2%-5% CH₃OH/CH₂Cl₂) gave
30 the title compound as a pale yellow foam (55%): ¹H NMR (CDCl₃) δ 8.22 (br s, 1H), 7.48 (s, 1H), 7.20 (d, J=8.3 Hz, 1H), 7.04 (d, J=8.3 Hz, 1H), 5.27 (d, J=16.6 Hz, 1H), 5.25 (br s, 1H), 3.92 (d, J=5.4 Hz, 2H), 3.82 (m, 1H), 3.80 (d, J=16.6 Hz, 1H), 3.71 (s, 3H), 3.03 (s, 3H), 2.90 (m, 3H), 2.41 (dd, J=16.8, 5.5 Hz, 1H), 1.48 (s, 9H).

35

b) (±)-8-[(2-Aminoacetyl)amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-benzazepine-2-acetic acid

- Methyl (±)-8-[(2-[tert-butoxycarbonyl]aminoacetyl)amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetate was saponified according to the procedure of example 6(b). The resulting product was stirred in 1:1 CH₂Cl₂/TFA for 2 h, then was concentrated. Purification by chromatography (ODS, 10% CH₃CN/H₂O-0.1% TFA) followed by lyophilization gave the title compound as a white powder (76%): ¹H NMR (DMSO-d₆) δ 8.10 (br s, 3H), 7.43 (d, J=8.2 Hz, 1H), 7.39 (s, 1H), 7.10 (d, J=8.2 Hz, 1H), 5.26 (d, J=16.1 Hz, 1H), 3.91 (d, J=16.1 Hz, 1H), 3.76 (m, 3H), 3.03 (dd, J=16, 2 Hz, 1H), 2.88 (s, 3H), 2.65 (m, 2H), 2.34 (dd, J=16, 2 Hz, 1H); MS (ES) m/e 306 (M+H)⁺. Anal. Calcd for C₁₅H₁₉N₃O₄ · 1.5 C₂H₅F₃O₂ · 0.5 H₂O: C, 44.54; H, 4.47; N, 8.66. Found: C, 44.76; H, 4.47; N, 8.43.

15

Example 21

Preparation of (+/-)-8-[(3-aminopropanoyl)amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid

- 20 a) Methyl (±)-8-[[3-[(tert-butoxycarbonyl)amino]propanoyl]amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetate

Methyl (±)-8-amino-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetate was coupled with Boc-β-alanine according to the procedure of Example 5(b). Purification by chromatography on silica gel gave the title compound (87%) as a light yellow foam: ¹H NMR (CDCl₃) δ 8.20 (br s, 1H), 7.48 (s, 1H), 7.20 (d, J=8.3 Hz, 1H), 7.04 (d, J=8.3 Hz, 1H), 5.27 (d, J=16.6 Hz, 1H), 3.92 (m, 3H), 3.80 (d, J=16.6 Hz, 1H), 3.70 (s, 3H), 3.01 (s, 3H), 2.90 (m, 5H), 2.48 (dd, J=16.8, 5.5 Hz, 1H), 1.48 (s, 9H).

- 30 b) (±)-8-[(3-Aminopropanoyl)amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid

Methyl (±)-8-[[3-[(tert-butoxycarbonyl)amino]propanoyl]amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetate was deprotected according to the procedure of Example 20(b). Purification by recrystallization (CH₃OH/EtOAc) gave the title compound as a white solid (45%): ¹H NMR (DMSO-d₆) δ 7.47 (s, 1H), 7.37 (d, J=8.3 Hz, 1H), 7.03 (d, J=8.3 Hz, 1H), 5.23 (d, J=16.3 Hz, 1H), 3.85 (d, J=16.3 Hz, 1H), 3.70 (m, 1H), 2.98 (m, 2H), 2.87 (s, 3H),

2.65 (m, 3H), 2.30 (dd, J=16, 6.8 Hz, 1H); MS (ES) m/e 320 (M+H)⁺. Anal. Calcd for C₁₆H₂₁N₃O₄ · 0.5 C₂F₃HO₂ · 0.5 H₂O: C, 52.37; H, 5.95; N, 10.78. Found: C, 52.29; H, 5.97; N, 10.78.

5

Example 22Preparation of (±)-4-methyl-3-oxo-7-[[[(3-pyridinyl)methyl]amino]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

- 10 a) Methyl (±)-4-methyl-3-oxo-7-[[[(3-pyridinyl)methyl]amino]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate
- EDC (229.4 mg, 1.2 mmol) was added to a solution of methyl (±)-7-carboxy-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (292.3 mg, 1.0 mmol), 3-(aminomethyl)pyridine (0.122 mL, 1.2 mmol),
- 15 HOBT·H₂O (162.2 mg, 1.2 mmol), and diisopropylethylamine (0.35 mL, 2.0 mmol) in anhydrous DMF (5 mL) at RT. After 23.5 h, the reaction was concentrated on the rotavap (high vacuum), and the residue was partitioned between 10% aqueous Na₂CO₃ and CHCl₃. This caused a solid to precipitate. The layers were separated and the aqueous layer was extracted exhaustively with CHCl₃
- 20 until all solids had dissolved. The combined CHCl₃ layers were diluted with CHCl₃, washed with 10% Na₂CO₃, dried (MgSO₄), and concentrated. Crystallization of the residue from EtOAc gave impure title compound. The mother liquors were concentrated, combined with the recrystallized residue and chromatographed (silica gel, 10% MeOH/CHCl₃). The title compound was
- 25 obtained as a colorless solid (341 mg, 89%): TLC R_f 0.37 (10% MeOH/CHCl₃); ¹H NMR (250 MHz, CDCl₃) δ 8.58-8.68 (m, 1H), 8.47-8.57 (m, 1H), 7.72-7.81 (m, 1H), 7.54 (d, J=2.1 Hz, 1H), 7.45 (dd, J=8.5, 2.1 Hz, 1H), 7.31 (dd, J=7.7, 4.8 Hz, 1H), 6.44-6.63 (m, 2H), 5.45 (d, J=16.4 Hz, 1H), 5.04-5.15 (m, 1H), 4.64 (d, J=5.9 Hz, 2H), 4.55 (br d, J=4.5 Hz, 1H), 3.76 (d, J=16.4 Hz, 1H), 3.75 (s, 3H), 3.07 (s,
- 30 3H), 2.99 (dd, J=16.0, 6.7 Hz, 1H), 2.67 (dd, J=16.0, 6.4 Hz, 1H); MS (ES) m/e 383.2 (M+H)⁺.

b) (±)-4-Methyl-3-oxo-7-[[[(3-pyridinyl)methyl]amino]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid

- 35 A mixture of methyl (±)-4-methyl-3-oxo-7-[[[(3-pyridinyl)methyl]amino]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetate (161.7 mg, 0.42 mmol), 1.0 N NaOH (1.3 mL, 1.3 mmol), and MeOH (4.2 mL) was stirred at 40°C


- 47 -

for 3 h, then at RT overnight. The resulting solution was concentrated, and the residue was dissolved in H₂O and CH₃CN. The solution was acidified to pH 1 with TFA and was concentrated. The residue was crystallized from H₂O to afford the title compound as an off-white, crystalline solid (120.9 mg, 60%) : HPLC k' 5.7
5 (PRP-1 \otimes , 8% CH₃CN/H₂O-0.1% TFA); ¹H NMR (250 MHz, DMSO-d₆) δ 8.60-8.83 (m, 3H), 8.15 (br d, J=8.0 Hz, 1H), 7.75 (dd, J=7.9, 5.3 Hz, 1H), 7.49-7.60 (m, 2H), 6.56 (d, J=9.1 Hz, 1H), 6.38 (br s, 1H), 5.49 (d, J=16.5 Hz, 1H), 5.01-5.16 (m, 1H), 4.37-4.65 (m, 2H), 3.82 (d, J=16.5 Hz, 1H), 2.92 (s, 3H), 2.77 (dd, J=16.7, 9.0 Hz, 1H), 2.61 (dd, J=16.7, 4.9 Hz, 1 H, partially obscured by the residual solvent
10 signal); MS (ES) m/e 369.0 (M+H)⁺. Anal. Calcd for C₁₉H₂₀N₄O₄ · CF₃CO₂H: C, 52.29; H, 4.39; N, 11.61. Found: C, 52.45; H, 4.40; N, 11.56.

The above description fully discloses how to make and use the present invention. However, the present invention is not limited to the particular
15 embodiments described hereinabove, but includes all modifications thereof within the scope of the following claims. The various references to journals, patents and other publications which are cited herein comprises the state of the art and are incorporated herein by reference as though fully set forth.

What is claimed is:

1. A compound according to formula (I):



wherein

X-X' is NR¹-CH, NC(O)R³-CH, N=C, CR¹=C, CHR¹-CH, O-CH or S-CH;

R¹ is H, C₁₋₆ alkyl or benzyl;

10 R² is (CH₂)_nCO₂H;

R³ is H, C₁₋₆alkyl, Ar-C₀₋₆alkyl, Het-C₀₋₆alkyl, or C₃₋₆cycloalkyl-C₀₋₆alkyl;

R⁴ is W-U, Y-(CHR⁵)_m-U or Z-C(O);


R⁵ and R⁶ are independently chosen from H, C₁₋₆alkyl, Ar-C₀₋₆alkyl,

Het-C₀₋₆alkyl and C₃₋₆cycloalkyl-C₀₋₆alkyl;



15 m is 1 or 2;

n is 1 or 2;

U is NR¹C(O), C(O)NR¹, CH=CH, C≡C, CH₂-CH₂, O-CH₂, CH₂-O or CH₂OCONR¹;



R^a is H, OH, NO₂, N(R⁶)₂, CON(R⁶)₂, CH₂N(R⁶)₂, or R⁶HN-C(=NH);



R^d is H, N(R¹)₂, C₁₋₄alkyl, CON(R¹)₂, OH, OR¹, or Ar-C₀₋₄alkyl;

25 R^e is H, C₁₋₄alkyl, 2- or 3- pyridinyl, 1-, 2- or 3-piperidinyl, or 2- or 4-pyrimidinyl;

- 49 -


and pharmaceutically acceptable salts thereof,
provided that R^3 is not phenylethyl when R^4 is
(3-amidino)phenylaminocarbonyl and X-X' is NH-CH.

5 2. A compound according to claim 1 which has a binding affinity to the vitronectin receptor relative to the fibrinogen receptor of greater than 3:1.

3. A compound according to claim 1 wherein X-X' is NH-CH or CH₂-CH and n
is 1.

10

4. A compound according to claim 1 wherein U is NR¹CO.

5. A compound according to claim 1 wherein W is 

15 6. A compound according to claim 1 wherein Z is 1-piperazinyl or 1-piperidinyl.

7. A compound according to claim 1 wherein R⁴ is Y-(CH₂)_mNCH₃CO.

20 8. A compound according to claim 1 wherein R^e is H or substituted or unsubstituted phenyl, benzyl, pyridinyl, pyrimidinyl or piperidinyl.

9. A compound according to claim 1 wherein Y is OH or NHR⁶.

10. A compound selected from the group of:

25 (±)-7-[[[(6-Amino-2-pyridinyl)amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;

(±)-8-[[[(6-amino-2-pyridinyl)amino]carbonyl]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid;

30 (±)-7-[[[(6-amino-3-pyridinyl)amino]carbonyl]-3-oxo-4-(2-phenylethyl)-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;

(±)-1-Acetyl-7-[[[(6-amino-2-pyridinyl)amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;

(±)-2-methyl-3-oxo-8-[[[2-(pyridinyl)carbonyl]amino]-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid;

35 (±)-8-[(benzyloxycarbonyl)amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid;

- 50 -

- (±)-7-[[[3-(aminoiminomethyl)phenyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;
- (±)-7-[[[3-(aminocarbonyl)phenyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;
- 5 (±)-4-methyl-3-oxo-7-[1-(4-phenylpiperazinyl)carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;
- (±)-4-methyl-3-oxo-7-[(1-piperazinyl)carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;
- (±)-4-methyl-3-oxo-7-[1-[4-(1-piperidinyl)piperidinyl]carbonyl]-2,3,4,5-tetrahydro-10 1H-1,4-benzodiazepine-2-acetic acid;
- (±)-4-methyl-3-oxo-7-[1-[4-(2-pyridinyl)piperazinyl]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;
- (±)-4-methyl-3-oxo-7-[1-[4-(phenylmethyl)piperazinyl]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;
- 15 (±)-4-methyl-3-oxo-7-[1-[4-(2-pyrimidinyl)piperazinyl]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;
- 8-[[2S-amino-3-phenylpropanoyl]amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4RS-acetic acid;
- (±)-8-[[[(2-hydroxy-2-phenyl)ethyl]methylamino]carbonyl]-2-methyl-3-oxo-20 2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid;
- (±)-7-[[[N-(2-hydroxyethyl)-N-methyl]amino]carbonyl]-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;
- (±)-8-[[[(2-(2-pyridinylamino)acetyl)amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid;
- 25 (±)-4-methyl-3-oxo-7-[(2-phenylaminoethyl)amino]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid;
- (±)-8-[(2-aminoacetyl)amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1h-2-benzazepine-4-acetic acid;
- (+/-)-8-[(3-aminopropanoyl)amino]-2-methyl-3-oxo-2,3,4,5-tetrahydro-1H-2-benzazepine-4-acetic acid; and
- 30 (±)-4-methyl-3-oxo-7-[[[(3-pyridinyl)methyl]amino]carbonyl]-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-2-acetic acid.

11. A pharmaceutical composition comprising a compound according to claim 1
35 and a pharmaceutically acceptable carrier.

- 51 -

12. A method of inhibiting a vitronectin receptor comprising administering a compound according to claim 1.

5 13. A method of for treating osteoporosis, atherosclerosis, cancer or restenosis in a mammal comprising administering a compound according to claim 1 and a pharmaceutically acceptable carrier.

14. The use of a compound according to claim 1 in the manufacture of a medicament for osteoporosis, atherosclerosis, cancer or restenosis.

10

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/08146

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) :A61K 31/55; C07D 243/14 US CL :514/221; 540/513, 504 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 514/221; 540/513, 504 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAS ONLINE		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO, A, 93/00095 (SMITHKLINE BEECHAM CORP.) 07 January 1993, see pages 1-16, 34-38, and 116-124.	1-14
X, P	WO, A, 94/14776 (SMITHKLINE BEECHAM CORP.) 07 July 1994, see entire document.	1-14
X	Bioorganic and Medicinal Chemistry, Volume 2, Number 9, issued 1994, Bondinell et al, "Design of a potent and orally active nonpeptide platelet fibrinogen receptor (GPIIb/IIIa) antagonist", pages 897-908, see entire document.	1 & 3-10
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 22 SEPTEMBER 1995		Date of mailing of the international search report 13 OCT 1995
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer KING L. WONG Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORTInt. application No.
PCT/US95/08146**C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	Tetrahedron Letters, Volume 36, Number 3, issued 1995, Miller et al, "Synthesis of a 2-Benzazepine Analog of a Potent, Nonpeptide GPIIb/IIIa Antagonist", pages 373-376, see entire document.	1 & 3-10
X, P	Journal of Medicinal Chemistry, Volume 38, Number 1, issued January 1995, Ku et al, "Potent Non-peptide Fibrinogen Receptor Antagonists Which Present an Alternative Pharmacophore", pages 9-12, see pages 10-11.	1 & 3-10
X	Journal of American Chemical Society, Volume 115, Number 19, issued September 1993, Ku et al, "Direct Design of a Potent Non-Peptide Fibrinogen Receptor Antagonist Based on the Structure and Conformation of a Highly Constrained Cyclic RGD Peptide", pages 8861-8862, see entire document.	1 & 3-10
X	WO, A, 93/08174 (GENENTECH, INC.) 29 April 1993, see pages 1-3, 5-10, 13, 16-18, 118-339, and 403.	1-14

INTERNATIONAL SEARCH REPORT

Int. application No.
PCT/US95/08146

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

- I. Claims 1-14, drawn to compounds, compositions, and methods of use, wherein X-X' is NR1-CH, NC(O)R3-CH, or N=C.
- II. Claims 1-14, drawn to compounds, compositions, and methods of use, wherein X-X' is CR1=C or CHR1-CH.
- III. Claims 1-14, drawn to compounds, compositions, and methods of use, wherein X-X' is O-CH.
- IV. Claims 1-14, drawn to compounds, compositions, and methods of use, wherein X-X' is S-CH.

The inventions of Groups I-IV do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons. The special technical feature of the Group I invention is diazepinyl compounds. Group II invention's special technical feature is azepinyl compounds. Group III invention's special technical feature is oxazepinyl compounds. Group IV invention's special technical feature is thiazepinyl compounds. The special technical feature of one of the above Groups is not present in the other Groups. There is no significant structural element shared by the inventions of Groups I-IV and the compounds are not art-recognized equivalents. Therefore, unity of invention is lacking.